

Meropenem Ranbaxy

Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

1. Why am I using Meropenem Ranbaxy?

Meropenem Ranbaxy contains the active ingredient meropenem. Meropenem Ranbaxy is used to treat infections of the lungs (pneumonia or bronchitis), infection of the bladder or kidney (urinary tract infection), infection of tissues around the stomach or gut, infections of the ovaries, vagina or womb, skin infections, infections of the membranes surrounding the brain and spinal cord (meningitis), infections of the blood (septicaemia), high fever and significant reduction in white blood cells resulting decreased resistance to infection (febrile neutropenia).

For more information, see Section [1. Why am I using Meropenem Ranbaxy?](#) in the full CMI.

2. What should I know before I use Meropenem Ranbaxy?

Do not use if you have ever had an allergic reaction to meropenem or any of the ingredients listed at the end of the CMI.

Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.

For more information, see Section [2. What should I know before I use Meropenem Ranbaxy?](#) in the full CMI.

3. What if I am taking other medicines?

Some medicines may interfere with Meropenem Ranbaxy and affect how it works.

A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

4. How do I use Meropenem Ranbaxy?

The dose of Meropenem Ranbaxy is based on your condition, age and type and severity of infection.

Your doctor will decide how much Meropenem Ranbaxy to give you. The usual dose is 500 mg to 1g injected every 8 hours. If you have meningitis you may require more. If you have kidney problems, a smaller dose may be given. Children also require lower doses.

More instructions can be found in Section [4. How do I use Meropenem Ranbaxy?](#) in the full CMI.

5. What should I know while using Meropenem Ranbaxy?

Things you should do	<ul style="list-style-type: none">Remind any doctor, dentist or pharmacist you visit that you are using Meropenem RanbaxyCall you doctor straight away if you develop severe diarrhoea
Driving or using machines	<ul style="list-style-type: none">Meropenem may cause headache, tingling or prickling skin and convulsions/seizures (fits) in some people. If you have any of these symptoms, do not drive, operate machinery or do anything that could be dangerous.
Looking after your medicine	<ul style="list-style-type: none">Meropenem Ranbaxy is store below 25°C

For more information, see Section [5. What should I know while using Meropenem Ranbaxy?](#) in the full CMI.

6. Are there any side effects?

Common side effects include pain with redness and/or swelling at the site of injection, feeling sick (nausea), being sick (vomiting), loose stool (diarrhoea), skin problems such as rash or itching, headache.

For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

Meropenem Ranbaxy

Active ingredient(s): *meropenem*

Consumer Medicine Information (CMI)

This leaflet provides important information about using Meropenem Ranbaxy. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using Meropenem Ranbaxy.**

Where to find information in this leaflet:

- [1. Why am I using Meropenem Ranbaxy?](#)
- [2. What should I know before I use Meropenem Ranbaxy?](#)
- [3. What if I am taking other medicines?](#)
- [4. How do I use Meropenem Ranbaxy?](#)
- [5. What should I know while using Meropenem Ranbaxy?](#)
- [6. Are there any side effects?](#)
- [7. Product details](#)

1. Why am I using Meropenem Ranbaxy?

Meropenem Ranbaxy contains the active ingredient meropenem. Meropenem Ranbaxy belongs to a group of medicines called carbapenem antibiotics. These medicines fight bacteria that cause infections in the body. Meropenem will not work against viral (such as colds and flu) or fungal diseases.

Meropenem Ranbaxy may have been prescribed to you or your child (aged 3 months or above) for the treatment of the following bacterial infections:

- Infections of the lungs (pneumonia or bronchitis)
- Infection of the bladder or kidney (urinary tract infection)
- Infection of tissues around the stomach or gut
- Infections of the ovaries, vagina or womb
- Skin infections
- Infections of the membranes surrounding the brain and spinal cord (meningitis)
- Infections of the blood (septicaemia)
- High fever and significant reduction in white blood cells resulting decreased resistance to infection (febrile neutropenia).

Meropenem Ranbaxy is given by injection, usually in hospitals, and its available only with a doctor's prescription.

Meropenem Ranbaxy is not addictive.

2. What should I know before I use Meropenem Ranbaxy?

Warnings

When you must not be given Meropenem Ranbaxy:

- you are allergic to meropenem
- you are allergic to any of the ingredients listed at the end of this leaflet
- you are allergic to other similar antibiotics (e.g. other carbapenems like imipenem; or beta-lactam antibiotics like penicillin, amoxicillin, ampicillin, ceftriaxone, cefotaxime etc).

Symptoms of allergic reaction may include skin rash, itchiness or hives, shortness of breath, wheezing or difficulty breathing, swelling of the face, tongue, lips or other parts of the body.

- Always check the ingredients to make sure you can use this medicine.

Check with your doctor if you:

- have any other medical conditions especially the following:
 - liver problems-if you notice yellowing of the skin and eyes, itchy skin, dark-coloured urine or light-coloured stool tell your doctor. This may be a sign of liver problems which your doctor needs to check.
 - kidney problems, including dialysis
 - disease of the stomach or intestines (in particular inflammation of intestine)
 - You may develop a positive test (Coomb's test) which indicates the presence of antibodies that may destroy red blood cells. Your doctor will discuss this with you.

Your doctor will discuss with you the benefits and risks of taking it, if you:

- have allergies to any other medicines, including antibiotics (such as other carbapenems, penicillins, cephalosporins or monobactams).
- have allergies to any foods, preservatives or dyes.
- are on a controlled sodium diet.
 - Meropenem Ranbaxy contains sodium (90 mg per 1 g of meropenem).

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

Pregnancy and breastfeeding

Check with your doctor if you are pregnant or intend to become pregnant.

Do not breast-feed if you are being given this medicine.

The active ingredient in Meropenem Ranbaxy passes into the breast milk. Your doctor may discuss the risk and benefits if you need this medicine while you are breast feeding.

Children

Safety and effectiveness of Meropenem Ranbaxy in children younger than 3 months of age have not been established. It is not recommended for use in these children.

Ask your doctor if you have any questions about why this medicine has been prescribed for you or your child. Your doctor may have prescribed it for another reason.

3. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and Meropenem Ranbaxy may interfere with each other. These include:

- Probenecid (a medicine used to treat gout)
- Sodium valproate (a medicine used to treat fit/seizure or mania)

These medicines may be affected by Meropenem Ranbaxy, or may affect how well it works. You may need different amounts of your medicine, or you may need to take different medicines. Your doctor will advise you.

Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect Meropenem Ranbaxy.

4. How do I use Meropenem Ranbaxy?

How much will you be given

The dose of Meropenem Ranbaxy is based on your condition, age and type and severity of infection.

Your doctor will decide how much Meropenem Ranbaxy to give you. The usual dose is 500 mg to 1g injected every 8 hours. If you have meningitis you may require more. If you have kidney problems, a smaller dose may be given. Children also require lower doses.

Meropenem Ranbaxy should not be given to babies less than three months of age or babies and children with kidney problems.

How Meropenem Ranbaxy is given

Meropenem Ranbaxy should only be injected into a vein (intravenous injection). It must always be given by a doctor

or a nurse. It can be given either as a slow injection or as a slow drip over 15 to 30 minutes.

Your doctor will decide which is best for you.

How long you will be given Meropenem Ranbaxy

Meropenem Ranbaxy will be given as a number of injections over several days. Your doctor will decide how long you will need to have Meropenem Ranbaxy.

If you use too much Meropenem Ranbaxy

As Meropenem is given to you under the supervision of your doctor, it is very unlikely that you will receive too much.

However, if you experience any severe side effects after being given Meropenem, you may need urgent medical attention.

You should immediately:

- phone the Poisons Information Centre (by calling **13 11 26**), or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

You should do this even if there are no signs of discomfort or poisoning.

5. What should I know while using Meropenem Ranbaxy?

Things you should do

Call your doctor straight away if you:

- Develop severe diarrhoea, even if it occurs several weeks after you have been given Meropenem Ranbaxy.

You may have developed a serious bowel condition called antibiotic-associated colitis and you need urgent medical attention. Do not take any medicine for your diarrhoea without checking with your doctor first.

Remind any doctor, dentist or pharmacist you visit that you are using Meropenem Ranbaxy.

Driving or using machines

Be careful before you drive or use any machines or tools until you know how Meropenem Ranbaxy affects you.

Meropenem may cause headache, tingling or prickling skin and convulsions/seizures (fits) in some people. If you have any of these symptoms, do not drive, operate machinery or do anything that could be dangerous. Children should be careful when undertaking activities requiring attention such as riding bicycles or climbing.

If you are about to have any blood tests, tell your doctor that you are being treated with Meropenem Ranbaxy.

It may interfere with results of some tests.

Looking after your medicine

The hospital staff will keep Meropenem Ranbaxy in a safe place below 25°C. They will also make sure that the expiry date has not passed.

Getting rid of any unwanted medicine

The hospital staff will dispose of any unused Meropenem Ranbaxy.

6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

Less serious side effects

Less serious side effects	What to do
<ul style="list-style-type: none"> • pain with redness and/or swelling at the site of injection • feeling sick (nausea), being sick (vomiting) • loose stool (diarrhoea) • skin problems such as rash or itching • headache. • reduced levels of potassium in your blood (which can cause weakness, muscle cramps, tingling and heart rhythm disturbances). 	<p>Speak to your doctor if you have any of these less serious side effects and they worry you.</p>

These are common side effects of meropenem, and are usually mild and short-lived.

Serious side effects

Serious side effects	What to do
<ul style="list-style-type: none"> • severe diarrhoea, even if it occurs several weeks after you have been given Meropenem Ranbaxy • allergic reactions, including shortness of breath, wheezing or difficulty in breathing; 	<p>Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of</p>

<p>swelling of the face, lips, tongue or other parts of the body; skin rash, itchiness or hives.</p> <ul style="list-style-type: none"> • chest pain or discomfort • fast or irregular heartbeat; or changes in the rhythm or rate of the heartbeat • convulsions/ seizures (fit) • Liver problems. Yellowing of the skin and eyes, itchy skin, dark-coloured urine or light-coloured stool. • unexpected breathlessness and /or red/brown urine- this may indicate damage to your red blood cells • muscle discomfort such as swelling, weakness, tender or sore muscles. 	<p>these serious side effects.</p>
--	---

Serious side effects are rare.

Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people.

Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at www.tga.gov.au/reporting-problems. By reporting side effects, you can help provide more information on the safety of this medicine.

Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.

7. Product details

This medicine is only available with a doctor's prescription.

What Meropenem Ranbaxy contains

Active ingredient (main ingredient)	Meropenem (as trihydrate)
Other ingredients (inactive ingredients)	Sodium carbonate anhydrous
Potential allergens	-

Do not take this medicine if you are allergic to any of these ingredients.

What Meropenem Ranbaxy looks like

Meropenem Ranbaxy 500 mg/20 mL vial (Aust R 167025) and 1 g/vial (AUST R 167024) is a sterile white to pale yellow powder supplied in a glass vial. The hospital staff make it up into a solution before injection.

Who distributes Meropenem Ranbaxy

Sun Pharma ANZ Pty Ltd

12 Waterloo Road

Macquarie Park Sydney NSW 2113 Australia

Tel: 1800 726 229

Email: customerservice.aus@sunpharma.com

Douglas Pharmaceuticals Ltd

P O Box 45 027

Auckland 0651

New Zealand

Phone: (09) 835 0660

This leaflet was prepared in February 2026.