

# TRACRIUM for Injection

*atracurium besilate*

## Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

### 1. Why am I using TRACRIUM?

TRACRIUM contains the active ingredient atracurium besilate. TRACRIUM will be given to you or your child during surgery. It is used together with anesthetic medicines to relax the body's muscles.

For more information, see Section [1. Why am I using TRACRIUM?](#) in the full CMI.

### 2. What should I know before I use TRACRIUM?

Do not use if you have ever had an allergic reaction to atracurium besilate or any of the ingredients listed at the end of the CMI.

**Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.**

For more information, see Section [2. What should I know before I use TRACRIUM?](#) in the full CMI.

### 3. What if I am taking other medicines?

Some medicines may interfere with TRACRIUM and affect how it works.

A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

### 4. How do I use TRACRIUM?

More instructions can be found in Section [4. How do I use TRACRIUM?](#) in the full CMI.

TRACRIUM will be given into your vein as an injection or by a 'drip' (infusion). This will be done by an anaesthetist or a doctor with special training.

### 5. What should I know while using TRACRIUM?

<b>Driving or using machines</b>	<ul style="list-style-type: none"><li>If you are discharged early, following treatment with TRACRIUM or any other anaesthetic agent, do not drive or operate machinery</li></ul>
<b>Looking after your medicine</b>	<ul style="list-style-type: none"><li>TRACRIUM should be stored by the hospital's pharmacy between 2° and 8°C and protected from light. Do not freeze. Once removed from the fridge, the ampoules may be stored below 25°C, protected from light for up to 1 week only, provided the product is used before the printed expiry date. Thereafter discard the ampoules.</li></ul>

For more information, see Section [5. What should I know while using TRACRIUM?](#) in the full CMI.

### 6. Are there any side effects?

The most commonly reported side effects are skin flushing, lowering of blood pressure, changes in heart rate either increased or decreased, tight breathing, allergic reactions, skin rash, seizures (in certain patients), muscle weakness.

For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

# TRACRIUM

Active ingredient(s): *atracurium besilate*

## Consumer Medicine Information (CMI)

This leaflet provides important information about using TRACRIUM. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using TRACRIUM.**

Where to find information in this leaflet:

- [1. Why am I using TRACRIUM?](#)
- [2. What should I know before I use TRACRIUM?](#)
- [3. What if I am taking other medicines?](#)
- [4. How do I use TRACRIUM?](#)
- [5. What should I know while using TRACRIUM?](#)
- [6. Are there any side effects?](#)
- [7. Product details](#)

### 1. Why am I using TRACRIUM?

TRACRIUM contains the active ingredient **atracurium besilate**. Tracrium belongs to a group of medicines called neuromuscular blockers. TRACRIUM works by blocking the effects of one of the body's chemical messengers called acetylcholine. Acetylcholine is involved in muscle contraction.

**Atracurium besilate is a highly selective, competitive or non-depolarising neuromuscular blocking agent which is used as an adjunct to general anaesthesia to enable tracheal intubation to be performed and to relax skeletal muscles during surgery or controlled ventilation, and to facilitate mechanical ventilation in Intensive Care Unit (ICU) patients.**

### 2. What should I know before I use TRACRIUM?

Do not use TRACRIUM if:

- you are allergic to atracurium besilate, or any of the ingredients listed at the end of this leaflet.** Symptoms of an allergic reaction may be mild or severe. They usually include some or all of the following:  
wheezing, swelling of the lips/mouth, difficulty in breathing, hayfever, lumpy rash ("hives") or fainting.

**Check with your doctor if you:**

- have had any adverse reactions during an operation.

- have had any type of allergic reaction to muscle relaxants, or to any medicines used during an operation.
- are allergic to any other medicines or any other substance, such as foods, dyes or preservatives.
- if you or a relative have had previous difficulties with anaesthetics.
- you have or have ever had any of the following medical conditions:
  - asthma or any other breathing disorders
  - heart disease
  - kidney, liver disease or disease of the nervous system e.g. myaesthesia gravis

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section 6. [Are there any side effects?](#)

### Pregnancy and breastfeeding

Check with your doctor if you are pregnant or intend to become pregnant.

Talk to your doctor if you are breastfeeding or intend to breastfeed.

Your doctor will discuss the possible risks and benefits of being given TRACRIUM if you are pregnant or breastfeeding.

**If you have not told your doctor about any of the above, tell them before you are given TRACRIUM.**

### 3. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

TRACRIUM may be affected by other medicines such as:

- antibiotics
- drugs for the heart
- blood pressure tablets
- magnesium
- lithium
- other muscle blockers (e.g. hexamethonium) and anaesthetics.

**Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect TRACRIUM.**

## 4. How do I use TRACRIUM?

TRACRIUM can be given into a vein in two ways:

- as an injection, or
- as a slow infusion.

TRACRIUM will be administered by an anaesthetist or other doctor with special training. You will never be expected to give yourself this medication. The dosage will vary according to many factors such as your body weight and the type of operation you have.

## 5. What should I know while using TRACRIUM?

### Driving or using machines

**Be careful before you drive or use any machines or tools until you know how TRACRIUM affects you.**

If you are discharged early, following treatment with TRACRIUM or any other anaesthetic agents, do not drive or operate machinery.

### Looking after your medicine

TRACRIUM should be stored by the hospital's pharmacy between 2° and 8°C and protected from light. Do not freeze.

Once removed from the fridge, the ampoules may be stored below 25°C, protected from light for up to 1 week only, provided the product is used before the printed expiry date. Thereafter discard the ampoules.

## 6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

Check with your doctor as soon as possible if you have any problems after receiving TRACRIUM, even if you do not think the problems are connected with the medicine or are not listed in this leaflet.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

### Serious side effects

Serious side effects	What to do
<ul style="list-style-type: none"><li>• skin flushing</li><li>• lowering of blood pressure</li><li>• changes in heart rate either increased or decreased</li><li>• tight breathing</li><li>• allergic reactions</li><li>• skin rash</li><li>• seizures (in certain patients)</li><li>• muscle weakness</li></ul>	<p><b>Call your doctor or nurse straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.</b></p>

**Tell your doctor, nurse or pharmacist if you notice anything else that may be making you feel unwell.**

Other side effects not listed here may occur in some people.

There is no evidence that TRACRIUM is addictive.

### Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at [www.tga.gov.au/reporting-problems](http://www.tga.gov.au/reporting-problems). By reporting side effects, you can help provide more information on the safety of this medicine.

## 7. Product details

This medicine is only available with a doctor's prescription.

### What TRACRIUM contains

<b>Active ingredient (main ingredient)</b>	atracurium besilate
<b>Other ingredients (inactive ingredients)</b>	benzenesulfonic acid, water for injections
<b>Potential allergens</b>	None

**Do not take this medicine if you are allergic to any of these ingredients.**

### What TRACRIUM looks like

TRACRIUM is clear, colourless to slightly yellow solution supplied in glass ampoules and is available at a concentration of 10 mg/mL in two sizes of 2.5 mL and 5 mL.

TRACRIUM 2.5 mL ampoules: AUST R 11008

TRACRIUM 5 mL ampoules: AUST R 39289

### Who distributes TRACRIUM

Aspen Pharmacare  
Australia Pty Ltd  
34-36 Chandos Street  
St Leonards NSW 2065  
Australia

This leaflet was prepared in February 2026.