

BTC Atorvastatin

Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

1. Why am I using BTC Atorvastatin?

BTC Atorvastatin contains the active ingredient atorvastatin. BTC Atorvastatin is used to lower high cholesterol levels.

BTC Atorvastatin is also used to help reduce the risk of having a heart attack or stroke in people who have high blood pressure and coronary heart disease (CHD) or who are at risk of CHD.

For more information, see Section [1. Why am I using BTC Atorvastatin?](#) in the full CMI.

2. What should I know before I use BTC Atorvastatin?

Do not use if you have ever had an allergic reaction to BTC Atorvastatin or any of the ingredients listed at the end of the CMI.

Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.

For more information, see Section [2. What should I know before I use BTC Atorvastatin?](#) in the full CMI.

3. What if I am taking other medicines?

Some medicines may interfere with BTC Atorvastatin and affect how it works.

A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

4. How do I use BTC Atorvastatin?

- The usual dose of BTC Atorvastatin is between 10-80 mg taken once a day.
- Swallow the tablets whole with a full glass of water. Do not chew or crush the tablets.

More instructions can be found in Section [4. How do I use BTC Atorvastatin?](#) in the full CMI.

5. What should I know while using BTC Atorvastatin?

| | |
|------------------------------------|--|
| Things you should do | <ul style="list-style-type: none">• Remind any doctor, dentist or pharmacist you visit that you are using BTC Atorvastatin.• Keep all of your doctor's appointments so that your progress can be checked. |
| Things you should not do | <ul style="list-style-type: none">• Do not stop using this medicine suddenly.• Do not give BTC Atorvastatin to anyone else, even if they have the same condition as you. |
| Driving or using machines | <ul style="list-style-type: none">• Be careful driving or operating machinery until you know how BTC Atorvastatin affects you.• Atorvastatin generally does not cause any problems with your ability to drive a car or operate machinery. However, as with many other medicines, it may cause dizziness in some people. |
| Drinking alcohol | <ul style="list-style-type: none">• Avoid drinking large quantities of alcohol.• Drinking large quantities of alcohol while taking BTC Atorvastatin may increase your chance of liver problems. |
| Looking after your medicine | <ul style="list-style-type: none">• Keep BTC Atorvastatin in a cool dry place where the temperature stays below 25°C.• Keep your tablets where children cannot reach them. |

For more information, see Section [5. What should I know while using BTC Atorvastatin?](#) in the full CMI.

6. Are there any side effects?

Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects. Tell your doctor or pharmacist if you notice any of the following and they worry you: muscle and joint pain, muscle weakness, especially in the forearms, thighs, hips, shoulders, neck, and back, difficulty climbing stairs or standing up from a chair, difficulty lifting arms over the head.

This is not a complete list of all possible side effects.

For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

BTC Atorvastatin

Active ingredient(s): *atorvastatin (as calcium trihydrate)*

Consumer Medicine Information (CMI)

This leaflet provides important information about using BTC Atorvastatin. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using BTC Atorvastatin.**

Where to find information in this leaflet:

- [1. Why am I using BTC Atorvastatin?](#)
- [2. What should I know before I use BTC Atorvastatin?](#)
- [3. What if I am taking other medicines?](#)
- [4. How do I use BTC Atorvastatin?](#)
- [5. What should I know while using BTC Atorvastatin?](#)
- [6. Are there any side effects?](#)
- [7. Product details](#)

1. Why am I using BTC Atorvastatin?

BTC Atorvastatin contains the active ingredient atorvastatin (as atorvastatin calcium trihydrate).

BTC Atorvastatin belongs to a group of medicines called HMG-CoA reductase inhibitors. It works by reducing the amount of cholesterol made by the liver. BTC Atorvastatin reduces the 'bad' cholesterol and can raise the 'good' cholesterol. BTC Atorvastatin also helps to protect you from a heart attack or stroke.

BTC Atorvastatin is used to help reduce the risk of having a heart attack or stroke in people who have high blood pressure and coronary heart disease (CHD) or who are at risk of CHD. Examples of risk factors for CHD include diabetes, a history of stroke, or small blood vessel disease.

Ask your doctor if you have any questions about why BTC Atorvastatin has been prescribed for you.

Your doctor may have prescribed it for another reason.

BTC Atorvastatin is not addictive.

What is cholesterol?

Everyone has cholesterol in their blood. It is a type of blood fat needed by the body for many things, such as building the cell lining, making bile acids (which help to digest food) and some hormones. However, too much cholesterol can be a problem.

Cholesterol is present in many foods and is also made in your body by the liver. If your body makes too much cholesterol or you have too much cholesterol in your diet, then your level becomes too high.

High cholesterol is more likely to occur with certain diseases or if you have a family history of high cholesterol.

There are different types of cholesterol. LDL, or low-density lipoprotein, is the 'bad' cholesterol that can block your blood vessels. HDL, or high-density lipoprotein, cholesterol is the 'good' cholesterol that is thought to remove the bad cholesterol from the blood vessels.

When you have high levels of 'bad' cholesterol in your blood, it may begin to 'stick' to the inside of your blood vessels instead of being carried to the parts of the body where it is needed. Over time, this can form hard areas, also called plaque, on the walls of your blood vessels, making it more difficult for the blood to flow. Sometimes, the plaque can detach from the vessel wall and float in the bloodstream; it can then reach a smaller vessel and completely block it. This blocking of your blood vessels can lead to several types of blood vessel disease, heart attack, angina and stroke.

There is another type of blood fat called triglyceride, which is a source of energy. However, high levels of triglyceride can be associated with a low level of 'good' cholesterol and may increase your risk of heart disease.

In some patients, BTC Atorvastatin is used to treat high cholesterol and high triglycerides together.

In most people, there are no symptoms of abnormal cholesterol or triglyceride levels. Your doctor can measure your levels with a simple blood test.

2. What should I know before I use BTC Atorvastatin?

Warnings

Do not use BTC Atorvastatin if:

- you are allergic to atorvastatin, or any of the ingredients listed at the end of this leaflet. Some of the symptoms of an allergic reaction may include
 - shortness of breath
 - wheezing or difficulty breathing
 - swelling of the face, lips, tongue or other parts of the body
 - rash, itching or hives on the skin.

Always check the ingredients to make sure you can use this medicine.

- you have active liver disease
- you are pregnant or intend to become pregnant
- you are taking the antibiotic fusidic acid hemihydrate which is used to treat infections
- you are taking the antiviral glecaprevir/pibrentasvir for the treatment of Hepatitis C
- you have intolerance to some sugars. Contact your doctor before taking BTC Atorvastatin as it contains lactose.

Do not take this medicine after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.

If it has expired or is damaged, return it to your pharmacist for disposal.

Check with your doctor if you have any of the following medical conditions:

- liver problems
- kidney problems
- muscle pain, tenderness or weakness from other medicines used to treat high cholesterol or triglycerides
- vision problems, eye muscle weakness and drooping eyelid
- a type of stroke called haemorrhagic stroke, or a type of stroke called a lacunar stroke.
If you have had one of these strokes before, this medicine may increase the risk of you having another haemorrhagic stroke
- breathing problems.

Your doctor will ask you to have your liver function tested before you start to take BTC Atorvastatin.

Tell your doctor if you drink alcohol regularly.

Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

Pregnancy and breastfeeding

Do not take this medicine if you are pregnant intend to become pregnant.

If you are a woman of child-bearing age and are taking this medicine, use a proven method of birth control to avoid pregnancy.

The medicine may affect your developing baby if you take it during pregnancy.

Do not breastfeed if you are taking this medicine.

The active ingredient in BTC Atorvastatin may pass into breast milk and affect your baby.

3. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines may be affected by BTC Atorvastatin or may affect how well it works. You may need different amounts of your medicines, or you may need to take different medicines. Your doctor will advise you.

Tell your doctor or pharmacist if you are taking any of the following:

- digoxin, a medicine used to treat some heart problems
- diltiazem, a medicine used to treat angina
- other medicines used to treat high cholesterol or triglycerides (fats)

- antacids, medicines used to treat reflux or ulcers
- the antibiotics erythromycin, clarithromycin, rifampicin, daptomycin or fusidic acid hemihydrate
- phenytoin, a medicine used to treat epilepsy (seizures)
- oral contraceptives for birth control
- ciclosporin, a medicine used to suppress the immune system
- some medicines used to treat fungal infections, such as Itraconazole or ketoconazole
- protease inhibitors for the treatment of HIV infection and/or Hepatitis C, such as efavirenz, fosamprenavir, ritonavir, boceprevir, telaprevir, tipranavir/ritonavir, elbasvir/grazoprevir and simeprevir
- HCV non structural protein 5A (NS5A)/5B (NS5B) inhibitors such as daclatasvir and ledipasvir
- letermovir
- spironolactone, a medicine used to treat high blood pressure and certain types of swelling
- vitamin B3
- colchicine, a medicine used to treat a disease with painful, swollen joints caused by uric acid crystals
- ticagrelor, a medicine used to prevent blood clots.

Your doctor and pharmacist have more information on medicines to be careful with or to avoid while taking BTC Atorvastatin.

Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect BTC Atorvastatin.

4. How do I use BTC Atorvastatin?

How much to take

- Follow the instructions provided and use BTC Atorvastatin until your doctor tells you to stop.
- Your doctor will tell you how many tablets you need to take each day. This may depend on your condition and whether or not you are taking any other medicines.
- The usual dose of BTC Atorvastatin is between 10-80 mg taken once a day.

When to take BTC Atorvastatin

- BTC Atorvastatin can be taken at any time of the day. However, your dose of BTC Atorvastatin should be taken at about the same time each day.
- Taking it at the same time each day will have the best effect. It will also help you remember when to take it.
- BTC Atorvastatin can be taken with or without food.

How to take BTC Atorvastatin

- Swallow the tablets whole with a glass of water. Do not crush or chew the tablets.
- Your doctor will discuss with you the need to be on a diet while you are taking BTC Atorvastatin. Follow your agreed diet plan carefully.

If you do not understand the instructions, ask your doctor or pharmacist for help.

How to long to take BTC Atorvastatin

- Take BTC Atorvastatin every day and continue taking it for as long as your doctor tells you.
- BTC Atorvastatin helps to lower your cholesterol, but it does not cure your condition. It is important to keep taking your medicine even if you feel well.
- If you stop taking BTC Atorvastatin, your cholesterol levels may rise again.

If you forget to use BTC Atorvastatin

BTC Atorvastatin should be used regularly at the same time each day.

If you miss your dose at the usual time:

- if it is less than 12 hours before your next dose, skip the dose you missed and take your next dose when you are meant to.
- otherwise, take it as soon as you remember, and then go back to taking your tablet as you would normally.

Do not take a double dose to make up for the dose that you missed.

This may increase the chance of you getting an unwanted side effect.

If you are not sure what to do, talk to your doctor or pharmacist.

If you use too much BTC Atorvastatin

If you think that you have used too much BTC Atorvastatin, you may need urgent medical attention.

You should immediately:

- phone the Poisons Information Centre (by calling 13 11 26), or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

You should do this even if there are no signs of discomfort or poisoning.

5. What should I know while using BTC Atorvastatin?

Things you should do

Keep all of your doctor's appointments so that your progress can be checked.

Your doctor will ask you to have your liver function tested from time to time while you are taking BTC Atorvastatin to make sure the medicine is working and to prevent unwanted side effects.

Your cholesterol and triglyceride levels also need to be checked regularly while you are taking this medicine.

A regular blood test to check an enzyme called creatine kinase (CK) may be needed. CK in your blood can rise after muscle injury which can be caused by medicines used to treat cholesterol or triglycerides, such as BTC Atorvastatin.

If you are about to be started on any new medicine, remind your doctor and pharmacist that you are taking BTC Atorvastatin.

Call your doctor straight away if you:

- become pregnant while you are taking BTC Atorvastatin. If you become pregnant while taking BTC Atorvastatin, stop taking it and contact your doctor immediately.

Remind any doctor, dentist or pharmacist you visit that you are using BTC Atorvastatin.

Things you should not do

- Do not stop using this medicine suddenly.
- Do not give BTC Atorvastatin to anyone else, even if they have the same condition as you.
- Do not take BTC Atorvastatin to treat any other complaints unless your doctor tells you to.

Things that would be helpful for your condition

Some self-help measures suggested below may assist your condition. Your doctor or pharmacist can give you more information about these measures.

- **weight:** While you are taking BTC Atorvastatin, you need to follow a diet plan agreed to with your doctor. This may include measures to lose some weight.
- **exercise:** Regular exercise can help lower your cholesterol levels. It is important not to overdo it. Before commencing regular exercise, you should consult your doctor who will suggest the most suitable exercise for you. If you experience any discomfort when exercising, see your doctor.
- **reduce alcohol:** Excessive alcohol intake can raise your cholesterol levels or affect your liver function, which could increase the chance of you getting unwanted side effects. Your doctor may discuss with you whether you should reduce the amount of alcohol you drink.
- **avoid smoking:** Smoking increases the risk of you suffering from heart problems. Your doctor may advise you to stop smoking.

Driving or using machines

Be careful before driving or operating machinery until you know how BTC Atorvastatin affects you.

Atorvastatin generally does not cause any problems with your ability to drive a car or operate machinery. However, as with many other medicines, it may cause dizziness in some people.

If you feel dizzy, do not drive, operate machinery or do anything else that could be dangerous.

Drinking alcohol

Avoid drinking large quantities of alcohol.

Drinking large quantities of alcohol while taking BTC Atorvastatin may increase your chance of getting liver problems.

Tell your doctor if you drink alcohol regularly.

Drinking grapefruit juice

Avoid drinking large quantities of grapefruit juice.

Grapefruit juice contains one or more components that alter the metabolism of some medicines, including atorvastatin.

Drinking very large quantities (over 1.2 liters) of grapefruit juice each day while taking atorvastatin increases your chance of getting side effects.

Looking after your medicine

- Keep your tablets in the blister pack until it is time to take them. If you take the tablets out of the blister pack they may not keep well.
Keep BTC Atorvastatin in a cool dry place where the temperature stays below 25°C.

Store it in a cool dry place away from moisture, heat or sunlight; for example, do not store it:

- in the bathroom or near a sink, or
- in the car or on window sills.

Keep it where young children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Getting rid of any unwanted medicine

If you no longer need to use this medicine or it is out of date, take it to any pharmacy for safe disposal.

Do not use this medicine after the expiry date, or if the packaging is torn or show signs of tampering.

6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

Less serious side effects

| Less serious side effects | What to do |
|--|--|
| Muscle-joint related <ul style="list-style-type: none"> muscle and joint pain, muscle weakness especially in the forearms, thighs, hips, shoulders, neck and back difficulty climbing stairs or standing up from a chair difficulty lifting arms over the head falling or difficulty getting up from a fall | Speak to your doctor if you have any of these less serious side effects and they worry you. These side effects are the more common side effects of your medicine. They are usually |
| Gut or Gastrointestinal related <ul style="list-style-type: none"> constipation, diarrhoea | |

| | |
|--|-----------------------|
| <ul style="list-style-type: none"> stomach or belly pain, nausea (feeling sick) heartburn, indigestion or wind | mils and short-lived. |
| General disorders <ul style="list-style-type: none"> headache | |
| Others <ul style="list-style-type: none"> urine infection stuffy or runny nose nose bleeds | |

Serious side effects

| Serious side effects | What to do |
|--|---|
| Liver related <ul style="list-style-type: none"> yellowing of skin and eyes and dark coloured urine | Call your doctor straight away if you notice any of these serious side effects. These are serious side effects that may require medical attention. Serious side effects are rare. |
| Kidney related <ul style="list-style-type: none"> feeling weak and tired, excessively thirsty and passing more urine | |
| Lungs related <ul style="list-style-type: none"> problems with breathing, including shortness of breath, persistent cough and fever | |
| Eye related <ul style="list-style-type: none"> vision problems, eye muscle weakness and drooping eyelids | |
| Others <ul style="list-style-type: none"> depression red spots or lumps on the skin, which could be signs of vasculitis (narrowing or blockage of blood vessels) skin rashes | |

Very serious side effects

| Very serious side effects | What to do |
|--|--|
| Allergy related <ul style="list-style-type: none"> symptoms of allergy such as skin rash, itching, swelling of the face, lips, mouth, tongue, throat or neck which may cause difficulty in swallowing or breathing | Go straight to the Emergency Department at your nearest hospital if you notice any of these very serious side effects. This list includes very serious side effects. You |
| Heart related <ul style="list-style-type: none"> chest pain | |
| Muscles related <ul style="list-style-type: none"> unexpected muscle pain, tenderness or weakness not | |

| | |
|--|--|
| <p>caused by exercise, particularly if you also feel unwell or have a fever</p> <p>Others</p> <ul style="list-style-type: none"> sudden severe headache, which may be accompanied by nausea, vomiting, loss of sensation, tingling in any part of the body or ringing in the ears severe blisters and bleeding of the lips, eyes, mouth, nose or genitals | <p>may need urgent medical attention or hospitalization.</p> |
|--|--|

Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people.

Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at www.tga.gov.au/reporting-problems. By reporting side effects, you can help provide more information on the safety of this medicine.

Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.

7. Product details

This medicine is only available with a doctor's prescription.

What BTC Atorvastatin contains

| | |
|---|---|
| Active ingredient (main ingredient) | atorvastatin calcium trihydrate |
| Other ingredients (inactive ingredients) | <ul style="list-style-type: none"> lactose monohydrate microcrystalline cellulose calcium carbonate croscarmellose sodium hydroxypropylcellulose polysorbate 80 magnesium stearate hypromellose Macrogol 8000 titanium dioxide purified talc Antifoam AF Emulsion Q7-2587 Euphorbia antisiphilitica leaf wax (only 10 mg, 20 mg and 40 mg) |
| Potential allergens | <ul style="list-style-type: none"> Sugars as lactose milk or milk products <p>BTC Atorvastatin tablets do not contain gluten, sucrose or glucose.</p> |

Do not take this medicine if you are allergic to any of these ingredients.

What BTC Atorvastatin looks like

BTC Atorvastatin 10 mg tablets – white to off-white, elliptical, film-coated tablets debossed with 'AS10' on one side and plain on the other side (Aust R 401445).

BTC Atorvastatin 20 mg tablets – white to off-white, elliptical, film-coated tablets debossed with 'AS20' on one side and plain on the other side (Aust R 401447).

BTC Atorvastatin 40 mg tablets – white to off-white, elliptical, film-coated tablets debossed with 'AS40' on one side and plain on the other side (Aust R 401449).

BTC Atorvastatin 80 mg tablets - white to off-white, elliptical, film-coated tablets debossed with 'AS80' on one side and plain on the other side (Aust R 401453).

Pack size of 30 tablets.

Who distributes BTC Atorvastatin

Sandoz Pty Ltd
 100 Pacific Highway
 North Sydney, NSW 2060
 Australia
 Tel 1800 726 369

This leaflet was prepared in March 2026.