

# CLONAC

## Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about taking this medicine, speak to your doctor or pharmacist.

### 1. Why am I taking CLONAC?

CLONAC contains the active ingredient diclofenac sodium. CLONAC is taken to treat pain and reduce inflammation (swelling and redness) that may occur in different types of arthritis including rheumatoid arthritis and osteoarthritis, other painful conditions where swelling is a problem such as back pain, rheumatism, muscle strains, sprains and tendonitis (e.g. tennis elbow), menstrual cramps (period pain). For more information, see Section [1. Why am I taking CLONAC?](#) in the full CMI.

### 2. What should I know before I take CLONAC?

Do not take CLONAC if you have ever had an allergic reaction to diclofenac sodium or any of the ingredients listed at the end of this leaflet. Do not take if you have had a stomach or intestinal ulcer, bleeding from the stomach or bowel (symptoms of which may include blood in your stools or black stools), severe kidney or liver problems, severe heart failure, heart bypass surgery.

Do not take CLONAC during the first 6 months of pregnancy, except on doctor's advice. Do not take CLONAC during the last three months of pregnancy. Use of this medicine during the last 3 months of pregnancy may affect your baby and may delay labour and birth. Use of non-aspirin NSAIDs can increase the risk of miscarriage, particularly when taken close to the time of conception. Tell your doctor if you are breastfeeding or wish to breastfeed. Like most NSAID medicines, CLONAC is not recommended for use during breastfeeding because CLONAC passes into breast milk and may affect your baby.

For more information, see Section [2. What should I know before I take CLONAC?](#) in the full CMI.

### 3. What if I am taking other medicines?

Some medicines may interfere with CLONAC and affect how it works.

A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

### 4. How do I take CLONAC?

- CLONAC tablets are usually taken in 2 or 3 doses during the day.
- Swallow the tablets whole with a full glass of water or other liquid. Do not chew them.

More instructions can be found in Section [4. How do I take CLONAC?](#) in the full CMI.

### 5. What should I know while taking CLONAC?

|                                  |   |
|----------------------------------|---|
| <b>Things you should do</b>      | <ul style="list-style-type: none"><li>• If you become pregnant while taking CLONAC, tell your doctor immediately. Your doctor can discuss with you the risk of taking it while you are pregnant.</li><li>• Ensure to keep all your doctor's appointments so that your progress can be checked.</li><li>• If you plan to have surgery, including dental surgery, tell your doctor or dentist that you are taking CLONAC.</li><li>• If you get an infection while taking CLONAC, tell your doctor.</li><li>• If you are about to be started on any new medicine, remind your doctor and pharmacist that you are taking CLONAC.</li></ul>  |
| <b>Things you should not do</b>  | <ul style="list-style-type: none"><li>• Do not take any of the following medicines while taking CLONAC without first telling your doctor: aspirin (also called ASA or acetylsalicylic acid), other salicylates, other medicines containing diclofenac (e.g. Voltaren tablets, Voltaren Emulgel), ibuprofen and any other NSAID medicines. If you take these medicines together with CLONAC, they may cause unwanted effects.</li><li>• If you need to take medicine for headache or fever, it is usually okay to take paracetamol. If you are not sure, your doctor or pharmacist can advise you.</li><li>• Do not stop any other forms of treatment for arthritis that your doctor has told you to follow.</li><li>• Do not take to treat any other conditions unless your doctor tells you to.</li><li>• Do not give your medicine to anyone else, even if they have the same condition as you.</li></ul> |
| <b>Driving or using machines</b> | <ul style="list-style-type: none"><li>• Be careful driving, operating machinery or doing jobs that require you to be alert until you know how CLONAC affects you. CLONAC may cause dizziness, drowsiness, spinning sensation (vertigo) or blurred vision in some people. If you have any of these symptoms, do not drive or do anything else that could be dangerous.</li></ul>   |

**Looking after  
your medicine**

- Keep your tablets in a cool dry place where the temperature stays below 25°C.

For more information, see Section [5. What should I know while taking CLONAC?](#) in the full CMI.

**6. Are there any side effects?**

There are a number of side effects associated with CLONAC. It is important to be aware of them so that you can identify any symptoms if they occur. For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

# CLONAC

Active ingredient: *diclofenac sodium*

## Consumer Medicine Information (CMI)

This leaflet provides important information about taking CLONAC. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about taking CLONAC.**

Where to find information in this leaflet:

- [1. Why am I taking CLONAC?](#)
- [2. What should I know before I take CLONAC?](#)
- [3. What if I am taking other medicines?](#)
- [4. How do I take CLONAC?](#)
- [5. What should I know while taking CLONAC?](#)
- [6. Are there any side effects?](#)
- [7. Product details](#)

## 1. Why am I taking CLONAC?

**CLONAC contains the active ingredient diclofenac sodium.** CLONAC belongs to a family of medicines called non-steroidal anti-inflammatory drugs (NSAIDs). These medicines can relieve the symptoms of pain and inflammation, but they will not cure your condition.

CLONAC is taken to treat pain and reduce inflammation (swelling and redness) that may occur in the following:

- Different types of arthritis including rheumatoid arthritis including rheumatoid and osteoarthritis.
- Other painful conditions where swelling is a problem such as back pain, rheumatism, muscle strains, sprains and tendonitis (e.g. tennis elbow).
- Menstrual cramps (period pain).

Your doctor may have prescribed this medicine for another reason.

Ask your doctor if you have any questions about why it has been prescribed for you. Your doctor may have prescribed it for another purpose.

CLONAC is only available with a doctor's prescription.

This medicine is not addictive.

## 2. What should I know before I take CLONAC?

### Warnings

**Do not take CLONAC if:**

- you are allergic (hypersensitive) to:
  - diclofenac sodium, or any of the ingredients listed at the end of this leaflet.
  - other medicines containing diclofenac sodium.
  - aspirin.
  - ibuprofen.

- any other NSAID medicines.

**If you are not sure if you are taking any of the above medicines, ask your doctor or pharmacist.**

Some of the symptoms of an allergic reaction may include:

- Shortness of breath.
- Wheezing or difficulty breathing.
- Swelling of the face, lips, tongue, throat, and/or extremities (signs of angioedema).
- Rash, itching or hives on the skin.

Many medicines used to treat headache, period pain and other aches and pains contain aspirin or NSAID medicines. If you are allergic to aspirin or NSAID medicines and take CLONAC, these symptoms may be severe.

**Do not take CLONAC if you have had any of the following medical conditions:**

- A stomach or intestinal ulcer.
- Bleeding from the stomach or bowel (symptoms of which may include blood in your stools or black stools).
- Severe kidney or liver problems.
- Severe heart failure.
- Heart bypass surgery.

**Do not take CLONAC during the first 6 months of pregnancy, except on doctor's advice. Do not take during the last three months of pregnancy.** Use of this medicine during the last 3 months of pregnancy may affect your baby and may delay labour and birth.

Use of non-aspirin NSAIDs can increase the risk of miscarriage, particularly when taken close to the time of conception.

Do not take CLONAC if the expiry date printed on the carton has passed.

Do not take CLONAC if the packaging is torn or shows signs of tampering.

Do not give CLONAC to children. CLONAC is not recommended for use in children, there is not enough information on its use in this age group.

**Tell your doctor if you have any medical conditions, especially the following:**

- Established disease of the heart or blood vessels (also called cardiovascular disease, including uncontrolled high blood pressure, congestive heart failure, established ischemic heart disease, peripheral arterial disease or atherosclerotic cardiovascular disease), as treatment with CLONAC is generally not recommended.
- Established cardiovascular disease (see above) or significant risk factors such as high blood pressure, abnormally high levels of fat (cholesterol, triglycerides) in your blood, diabetes, or if you smoke, and your doctor decides to prescribe CLONAC, you

must not increase the dose above 100 mg per day if you are treated for more than 4 weeks.

- A past history of ulcers (stomach or intestinal).
- Gastrointestinal problems such as stomach ulcer, bleeding or black stools, or have experienced stomach discomfort or heartburn after taking anti-inflammatory medicines in the past.
- Diseases of the bowel or inflammation of the intestinal tract (Crohn's disease) or colon (ulcerative colitis).
- Past history of haemorrhoids (piles) or irritation of the rectum (back passage).
- Liver or kidney problems.
- A rare liver condition called porphyria.
- Bleeding disorders or other blood disorders (e.g. anaemia).
- Asthma or any other chronic lung disease that causes difficulty in breathing.
- Hay fever (seasonal allergic rhinitis).
- Repeated chest infections.
- Polyps in the nose.
- Diabetes.
- Dehydration (e.g. by sickness, diarrhoea, before or after recent major surgery).
- Swollen feet.

Your doctor may want to take special care if you have any of these conditions.

It is generally important to take the lowest dose that relieves your pain and/ or swelling and for the shortest time possible in order to keep your risk for heart problems or high blood pressure as small as possible.

#### **Tell your doctor if you currently have an infection.**

If you take CLONAC while you have an infection, some of the signs of the infection such as pain, fever, swelling and redness may be hidden. You may think, mistakenly, that you are better or that the infection is not serious.

#### **Tell your doctor if you are lactose intolerant.**

CLONAC tablets contain lactose.

#### **Tell your doctor if you are allergic to any other medicines, foods, dyes or preservatives.**

Your doctor will want to know if you are prone to allergies, especially if you get skin reaction with redness, itching or rash.

If you have not told your doctor about any of the above, tell your doctor before taking CLONAC.

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

### **Pregnancy and breastfeeding**

#### **Tell your doctor if you are pregnant or plan to become pregnant.**

There is not enough information to recommend CLONAC during the first 6 months of pregnancy and it must not be used during the last 3 months. This medicine may also

reduce fertility and affect your chances of becoming pregnant. Your doctor can discuss with you the risks and benefits involved.

**Tell your doctor if you are breastfeeding or wish to breastfeed.** Like most NSAID medicines, CLONAC is not recommended for use during breastfeeding because CLONAC passes into breast milk and may affect your baby.

### **3. What if I am taking other MEDICINES?**

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

You may need to take different amounts of your medicine or to take different medicines while you are taking CLONAC. Your doctor and pharmacist have more information.

#### **Some medicines may interfere with CLONAC and affect how it works:**

- Other anti-inflammatory medicines, e.g. aspirin, salicylates or ibuprofen.
- Warfarin or other "blood thinners" (medicines used to prevent blood clotting).
- Digoxin, a medicine for heart problems.
- Lithium or selective serotonin - reuptake inhibitors (SSRIs), a medicine used to treat some types of depression.
- Diuretics, medicines used to increase the amount of urine.
- ACE inhibitors or beta-blockers, medicines used to treat high blood pressure, heart conditions, glaucoma and migraine.
- Prednisone, cortisone, or other corticosteroids, medicines used to provide relief for inflamed areas of the body.
- Medicines (such as metformin) used to treat diabetes, except insulin.
- Methotrexate, a medicine used to treat arthritis and some cancers.
- Ciclosporin and tacrolimus, medicines used in patients who have received organ transplants.
- Trimethoprim used to treat urinary tract infections.
- Some medicines used to treat infection, quinolone antibacterials.
- Glucocorticoid medicines, used to treat arthritis.
- Sulfinpyrazone, a medicine used to treat gout.
- Voriconazole, a medicine used to treat fungal infections.
- Phenytoin, a medicine used to treat seizures.
- Rifampicin (an antibiotic medicine used to treat bacterial infections).

### **4. How do I take CLONAC?**

#### **How much to take**

##### **To Treat Arthritis or Other Painful Conditions**

The usual starting dose of CLONAC tablets is 75 mg to 150 mg each day.

After the early stages of treatment, it is usually possible to reduce the dose to 75 mg to 100 mg each day.

### **To Treat Menstrual Cramps (period pain)**

The tablets are usually taken during each period as soon as cramps begin and continued for a few days until the pain goes away.

The usual starting dose of CLONAC tablets is 50 mg to 100 mg each day, beginning as soon as cramps begin and continuing until the pain goes away, but for no longer than 3 days.

If necessary, the dose can be raised over several menstrual periods to a maximum of 200 mg each day.

Follow all directions given to you by your doctor and pharmacist carefully.

These instructions may differ from the information contained in this leaflet.

If you do not understand the instructions on the carton, ask your doctor or pharmacist for help.

There are different ways to take this medicine depending upon your condition. Your doctor will tell you exactly how many tablets to take.

**Elderly patients should take the minimum number of tablets that provides relief of symptoms.** Elderly patients may be more sensitive to the effects of CLONAC than other adults.

**Do not exceed the recommended dose.**

### **When to take CLONAC**

It is recommended to take CLONAC tablets before meals or on an empty stomach. If they upset your stomach, you can take them with food or immediately after food.

They will work more quickly if you take them on an empty stomach but they will still work if you have to take them with food to prevent stomach upset.

### **How to take CLONAC**

CLONAC tablets are usually taken in 2 or 3 doses during the day.

**Swallow the tablets whole with a full glass of water. Do not chew them.** The tablets have a special coating to stop them dissolving until they have passed through the stomach into the bowel. Chewing them would destroy the coating.

### **How long to take CLONAC**

Do not take CLONAC for longer than your doctor says.

If you are taking CLONAC for arthritis, it will not cure your disease but it should help control pain and inflammation. It usually begins to work within a few hours but several weeks may pass before you feel the full effects of the medicine.

### **If you forget to take CLONAC**

If it is almost time for your next dose (e.g. within 2 or 3 hours), skip the dose you missed and take your next dose when you are meant to.

Otherwise, take the missed dose as soon as you remember, and then go back to taking your tablets as you would normally.

Do not take a double dose to make up for the dose you missed. This may increase the chance of you getting unwanted side effects.

**Ask your doctor or pharmacist if you are not sure what to do.**

### **If you take too much CLONAC**

If you think that you have taken too much CLONAC, you may need urgent medical attention.

**You should immediately:**

- phone the Poisons Information Centre (by calling 13 11 26), or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

**You should do this even if there are no signs of discomfort or poisoning.**

If you take too much CLONAC, you may experience:

- Vomiting.
- Bleeding from the stomach or bowel.
- Diarrhoea.
- Dizziness.
- Ringing in the ears.
- Convulsions (fits).

## **5. What should I know while taking CLONAC?**

### **Things you should do**

**If you take CLONAC for more than a few weeks, you should make sure you visit your doctor for regular check-ups to ensure that you are not suffering from unnoticed undesirable side effects.**

**If you become pregnant while taking CLONAC, tell your doctor immediately.** Your doctor can discuss with you the risk of taking it while you are pregnant.

**Be sure to keep all your doctor's appointments so that your progress can be checked.**

Your doctor will periodically re-evaluate whether you should continue treatment, if you have established heart disease or significant risks for heart disease, especially in case you are treated for more than 4 weeks.

Your doctor may want to check your kidneys, liver and blood from time to time to help prevent unwanted side effects.

If, at any time while taking CLONAC you experience any signs or symptoms of problems with your heart or blood vessels such as chest pain, shortness of breath, weakness,

or slurring of speech, contact your doctor immediately. These may be signs of cardiovascular toxicity.

**If you plan to have surgery, including dental surgery, tell your doctor or dentist that you are taking CLONAC.** NSAID medicines can slow down blood clotting and affect kidney function.

**If you get an infection while taking CLONAC, tell your doctor.** This medicine may hide some of the signs of an infection (pain, fever, swelling, redness). You may think, mistakenly, that you are better or that the infection is not serious.

**If you are about to be started on any new medicine, remind your doctor and pharmacist that you are taking CLONAC.** Tell any other doctors, dentists and pharmacists who are treating you that you are taking CLONAC.

### Things you should not do

**Do not take any of the following medicines while you are taking CLONAC without first telling your doctor:**

- Aspirin (also called ASA or acetylsalicylic acid).
- Other salicylates.
- Other medicines containing diclofenac (e.g. Voltaren tablets, Voltaren Emulgel).
- Ibuprofen.
- Any other NSAID medicines.

If you take these medicines together with CLONAC, they may cause unwanted effects.

If you need to take medicine for headache or fever, it is usually okay to take paracetamol. If you are not sure, your doctor or pharmacist can advise you.

**Do not stop any other forms of treatment for arthritis that your doctor has told you to follow.** This medicine does not replace exercise or rest programs or the use of heat/cold treatments.

**Do not take CLONAC to treat any other conditions unless your doctor tells you to.**

**Do not give your medicine to anyone else, even if they have the same condition as you.**

### Driving or using machines

**Be careful driving, operating machinery or doing jobs that require you to be alert until you know how CLONAC affects you.** CLONAC may cause dizziness, drowsiness, spinning sensation (vertigo) or blurred vision in some people. If you have any of these symptoms, do not drive or do anything else that could be dangerous.

### Looking after your medicine

Follow the instructions on the carton on how to take care of your medicine properly.

Keep your tablets in the blister pack until it is time to take them. If you take the tablets out of the blister pack they may not keep well.

Store your tablets in a cool dry place where the temperature stays below 25°C. Store away from moisture, heat or sunlight; for example, do not store it:

- in the bathroom or near a sink, or
- in the car or on window sills.

**Keep it where young children cannot reach it.**

### Getting rid of any unwanted medicine

If you no longer need to take CLONAC or it is out of date, take it to any pharmacy for safe disposal.

Do not take CLONAC after the expiry date.

## 6. Are there any side effects?

**Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking CLONAC.**

Like all other medicines, CLONAC may have unwanted side effects in some people. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects.

**If you are over 65 years of age, you may have an increased chance of getting side effects. Report any side effects to your doctor.** As people grow older, they are more likely to get side effects from medicines.

Do not be alarmed by this list of possible side effects.

You may not experience any of them.

### Less serious side effects

| Less serious side effects  | What to do  |
|--|---|
| <ul style="list-style-type: none"><li>• Stomach upset including nausea (feeling sick), vomiting, indigestion, cramps, loss of appetite, wind.</li><li>• Heartburn or pain behind or below the breastbone (possible symptoms of an ulcer in the tube that carries food from the throat to the stomach).</li><li>• Stomach or abdominal pain.</li><li>• Constipation, diarrhoea.</li><li>• Sore mouth or tongue.</li><li>• Altered taste sensation.</li><li>• Headache.</li><li>• Dizziness, spinning sensation.</li><li>• Drowsiness, disorientation, forgetfulness.</li><li>• Feeling depressed, anxious or irritable.</li><li>• Strange or disturbing thoughts or moods.</li><li>• Shakiness, sleeplessness, nightmares.</li><li>• Tingling or numbness of the hands or feet.</li><li>• Feeling of fast or irregular heartbeat.</li></ul> | <p><b>Speak to your doctor if you have any of these less serious side effects and they worry you.</b></p> |

| Less serious side effects   | What to do  |
|---|---|
| <ul style="list-style-type: none"> <li>• Unusual weight gain or swelling of arms, hands, feet, ankles or legs due to fluid build-up.</li> <li>• Symptoms of sunburn (such as redness, itching, swelling, blistering of the lips, eyes, mouth, and/or skin) that happen more quickly than normal.</li> <li>• Skin inflammation with flaking or peeling.</li> <li>• Vision disorders (e.g. blurred or double vision). If symptoms of vision disorders occur during treatment with CLONAC, contact your doctor as an eye examination may be considered to exclude other causes.</li> <li>• Buzzing or ringing in the ears, difficulty hearing.</li> <li>• Hypertension (high blood pressure).</li> <li>• Hair loss or thinning.</li> </ul> |   |
| <ul style="list-style-type: none"> <li>• NSAIDs, including diclofenac, may be associated with increased risk of gastrointestinal anastomotic leak.</li> </ul>   | <p><b>Close medical surveillance and caution are recommended when taking CLONAC after gastrointestinal surgery.</b></p> |

### Serious side effects

| Serious side effects   | What to do   |
|--|--|
| <ul style="list-style-type: none"> <li>• Red or purple skin (possible signs of blood vessel inflammation).</li> <li>• Severe pain or tenderness in the stomach, vomiting blood or material that looks like coffee grounds, bleeding from the back passage, black sticky bowel motions (stools) or bloody diarrhoea (possible stomach problems).</li> <li>• Rash, skin rash with blisters, itching or hives on the skin; swelling of the face, lips, mouth, tongue, throat, or other part of the body which may cause difficulty to swallow, low blood pressure (hypotension), fainting, shortness of breath (possible allergic reaction).</li> <li>• Wheezing, troubled breathing, or feelings of tightness in the chest (signs of asthma).</li> </ul> | <p><b>Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.</b></p> |

| Serious side effects  | What to do |
|---|------------|
| <ul style="list-style-type: none"> <li>• Yellowing of the skin and/or eyes (signs of hepatitis/ liver failure).</li> <li>• Persistent nausea, loss of appetite, unusual tiredness, vomiting, pain in the upper right abdomen, dark urine or pale bowel motions (possible liver problems).</li> <li>• Constant "flu-like" symptoms including chills, fever, sore throat, aching joints, swollen glands, tiredness or lack of energy, bleeding or bruising more easily than normal (possible blood problems).</li> <li>• Painful red areas, large blisters, peeling of layers of skin, bleeding in the lips, eyes, mouth, nose or genitals, which may be accompanied by fever and chills, aching muscles and feeling generally unwell (possible serious skin reaction).</li> <li>• Signs of a possible effect on the brain, such as sudden and severe headache, stiff neck (signs of viral meningitis), severe nausea, dizziness, numbness, difficulty in speaking, paralysis (signs of cerebral attack), convulsions (fits).</li> <li>• Change in the colour or amount of urine passed, frequent need to urinate, burning feeling when passing urine, blood or excess protein in the urine (possible kidney disorders).</li> <li>• Sudden and oppressive chest pain may be a sign of myocardial infarction or heart attack.</li> <li>• Breathlessness, difficulty breathing when lying down, swelling of the feet or legs (signs of cardiac failure).</li> <li>• Coincidental occurrence of chest pain and allergic reactions (signs of Kounis syndrome).</li> </ul> |            |

**Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.**

Other side effects not listed here may occur in some people.

### Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at [www.tga.gov.au/reporting-problems](http://www.tga.gov.au/reporting-problems). By reporting side

effects, you can help provide more information on the safety of this medicine.

**Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.**

## 7. Product details

This medicine is only available with a doctor's prescription.

### What CLONAC contains

|   |   |
|---|---|
| <b>Active ingredient (main ingredient)</b>      | <ul style="list-style-type: none"><li>• Diclofenac sodium</li></ul>   |
| <b>Other ingredients (inactive ingredients)</b> | <ul style="list-style-type: none"><li>• Colloidal anhydrous silica.</li><li>• Microcrystalline cellulose.</li><li>• Lactose monohydrate.</li><li>• Magnesium stearate.</li><li>• Maize starch.</li><li>• Povidone.</li><li>• Hypromellose.</li><li>• Iron oxide yellow.</li><li>• Titanium dioxide.</li><li>• Sodium starch glycollate type A.</li><li>• PEG-40 hydrogenated castor oil.</li><li>• Methacrylic acid-ethyl acrylate copolymer (1:1).</li><li>• Triethyl citrate.</li><li>• Iron oxide red (50 mg tablet only).</li></ul> |
| <b>Potential allergens</b>                      | Contains sugars as lactose.   |

This medicine is gluten free.

**Do not take this medicine if you are allergic to any of these ingredients.**

### What CLONAC looks like

#### CLONAC 25 mg Tablets

Pale yellow, round, enteric coated tablets plain on both sides. AUST R 272878.

Blister pack containing 20 or 50 tablets\*.

#### CLONAC 50 mg Tablets

Pale brown, round, enteric coated tablets plain on both sides. AUST R 272874.

Blister pack containing 10, 20 or 50 tablets\*.

\*Not all pack size are available.

### Who distributes CLONAC

Arrotex Pharmaceuticals Pty Ltd  
15 - 17 Chapel Street,  
Cremorne, VIC 3121

<http://arrotex.com.au/>

This leaflet was prepared in December 2024.