

ROBINUL injection

Consumer Medicine Information (CMI) summary

The [full CMI on the next page](#) has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

1. Why am I using ROBINUL Injection?

ROBINUL injection contains the active ingredient glycopyrronium bromide.

ROBINUL injection is used before surgery to decrease the volume of secretions from your mouth, lungs and stomach. It can also be used either before or during surgery to maintain your heart's normal beating rhythm. ROBINUL injection is also used to counter the effects of some other medicines, which can slow your heartbeat or produce excessive secretions when used during surgery.

For more information, see Section [1. Why am I using ROBINUL injection?](#) in the full CMI.

2. What should I know before I use ROBINUL injection?

Do not use if you have ever had an allergic reaction to ROBINUL injection or any of the ingredients listed at the end of the CMI.

Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.

For more information, see Section [2. What should I know before I am given ROBINUL injection?](#) in the full CMI.

3. What if I am taking other medicines?

Some medicines may interfere with ROBINUL injection and affect how it works.

A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

4. How ROBINUL Injection is given?

- The dose of ROBINUL may be different for each person.
- The dose given depends on your condition, age, body weight and your doctor.
- Your doctor will decide the right dose for you.
- For children aged 1 month to 12 years, the usual dose depends on their bodyweight.

More instructions can be found in Section [4. How ROBINUL injection is given?](#) in the full CMI.

5. What should I know while using ROBINUL Injection?

Things you should do	<ul style="list-style-type: none">• Remind any doctor, dentist or pharmacist who is treating you that you have had ROBINUL injection.• Tell your doctor if you feel ROBINUL is not helping your condition.
Things you should be careful of	<ul style="list-style-type: none">• Be careful during warm weather and temperature, and/or with physical exercise after use of ROBINUL. It may reduce your ability to sweat and can therefore cause overheating.• Be careful if you are elderly, unwell or taking other medicines. Some people may experience side effects such as drowsiness, or blurred vision, which may increase the risk of a fall.
Driving or using machines	<ul style="list-style-type: none">• Do not drive or operate machinery until you know how ROBINUL injection affects you. ROBINUL may cause drowsiness or dizziness or blurred vision in some people and therefore may affect alertness.
Looking after your medicine	<ul style="list-style-type: none">• ROBINUL injection will be stored in the pharmacy or on the hospital ward. The injection is kept in a cool dry place where the temperature stays below 25°C.

For more information, see Section [5. What should I know when given ROBINUL injection?](#) in the full CMI.

6. Are there any side effects?

Tell your doctor or pharmacist as soon as possible if you do not feel well after having ROBINUL injection.

For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

ROBINUL injection

Active ingredient: *glycopyrronium bromide*

Consumer Medicine Information (CMI)

This leaflet provides important information about using ROBINUL injection.

You should also speak to your doctor or nurse if you would like further information or if you have any concerns or questions about using ROBINUL injection.

Where to find information in this leaflet:

- [1. Why am I using ROBINUL injection?](#)
- [2. What should I know before I am given ROBINUL injection?](#)
- [3. What if I am taking other medicines?](#)
- [4. How ROBINUL injection is given?](#)
- [5. What should I know when given ROBINUL injection?](#)
- [6. Are there any side effects?](#)
- [7. Product details](#)

1. Why am I using ROBINUL injection?

ROBINUL injection contains the active ingredient glycopyrronium bromide. It is used before surgery to decrease the volume of secretions from your mouth, lungs and stomach. It can also be used either before or during surgery to maintain your heart's normal beating rhythm.

ROBINUL is also used to counter the effects of some other medicines, which can slow your heartbeat or produce excessive secretions when used during surgery

ROBINUL works in the nervous system by blocking an enzyme, which is responsible for producing saliva in your mouth and secretions in your stomach and lungs. This enzyme also slows down the heart rate. When this enzyme is blocked, secretions dry up and the heart rate increases.

Ask your doctor if you have any questions about why ROBINUL injection has been prescribed for you.

Your doctor may have prescribed it for another purpose. ROBINUL injection is not addictive.

2. What should I know before I am given ROBINUL injection?

Warnings

You must not be given ROBINUL injection if:

- you are allergic to glycopyrronium bromide or any of the ingredients listed at the end of this leaflet.** Some of the symptoms of an allergic reaction may include rash, itching, hives on the skin, swelling of the

face, lips, tongue or other parts of the body, shortness of breath, wheezing or troubled breathing.

- the expiry date printed on the pack has passed.**
- the packaging is torn or shows signs of tampering.**

Check with your doctor if you:

- have or have had any other medical conditions, such as:**
 - asthma
 - glaucoma, which is an increased pressure in your eyes
 - diarrhoea
 - problems with your heart, kidney, or thyroid
 - difficulty urinating, or enlarged prostate
 - stomach or bowel problems
 - hiatus hernia that causes heartburn and belching
 - high blood pressure
 - myasthenia gravis, a muscle weakness disorder
 - nerve or brain disorder, brain damage, or Down's Syndrome.
- allergies to any other substances, such as foods, preservatives, dyes or latex rubber.**

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

Pregnancy and breastfeeding

Check with your doctor if you are pregnant or intend to become pregnant.

Your doctor will discuss the risks and benefits of receiving ROBINUL during pregnancy.

Talk to your doctor if you are breastfeeding or intend to breastfeed.

Medicines like ROBINUL may reduce your milk supply. It is not known if ROBINUL is found in breast milk.

3. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines may interfere with ROBINUL and affect how it works. These include:

- ritodrine hydrochloride
- medicines for Parkinson's disease
- phenothiazines, which are used to treat mental illness or severe nausea and vomiting
- antidepressants
- antihistamines
- narcotic pain killers such as pethidine

- medicines used to control heart rhythm
- digoxin used to treat heart failure
- thioxanthenes
- butyrophenones
- amantadine.

You may need to take different amounts of your medicine, or you may need to take different medicines.

Your medical practitioner has more information on medicines to be careful with or avoid while being treated with ROBINUL injection.

4. How ROBINUL injection is given?

How much is given

Your doctor will decide what dose you will receive. This depends on your condition, age and body weight.

- *When used before surgery:* the usual adult dose is 0.2 to 0.4 mg given by injection into either a vein or a muscle. You will get ROBINUL before the anaesthetic is given.
- *For use during surgery:* the usual adult dose is a single injection of 0.2 to 0.4 mg into a vein. Sometimes the dose may need to be repeated. Your doctor will decide if this needs to happen.
- *To counter the effects of other medicines:* the usual adult dose of ROBINUL is 0.2 mg per 1 mg neostigmine or the equivalent dose of pyridostigmine given by injection into a vein.
- *For children aged 1 month to 12 years:* the usual dose depends on their bodyweight.

How it is given

- This medicine is normally injected into a muscle or a vein by a doctor or a nurse in a hospital setting.
- If ROBINUL is given to counter the effects of other medicines, it may be given in the same syringe as the neostigmine or pyridostigmine.

How long is it given

ROBINUL injection is usually given as a single dose, but sometimes a dose must be repeated. Your doctor will decide how many doses of ROBINUL injection are needed.

5. What should I know after I am given ROBINUL injection?

Things you should do

Tell your doctor or nurse if:

- you have any questions about why this medicine has been given to you.
- you feel ROBINUL is not helping your condition.

Remind any doctor, dentist or nurse you visit that you have been given ROBINUL injection.

Things to be careful of

- **Be careful if you are elderly, unwell or taking other medicines.**

Some people may experience side effects such as drowsiness, or blurred vision, which may increase the risk of a fall.

- **Be careful during warm weather and temperature, and/or with physical exercise after use of ROBINUL.**

It may reduce your ability to sweat and can therefore cause overheating.

Driving or using machines

Make sure you know how you react to ROBINUL injection before you drive a car, operate machinery, or do anything else that could be dangerous if you are drowsy, dizzy, or not alert.

Looking after this medicine

ROBINUL injection will be stored in the pharmacy or on the hospital ward. The injection is kept in a cool dry place where the temperature stays below 25°C.

6. Are there any side effects?

Tell your doctor, pharmacist or nurse as soon as possible if you do not feel well while you are being treated with ROBINUL injection.

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

Less serious side effects

Less serious side effects	What to do
<ul style="list-style-type: none"> • fast, slow or irregular heart beats • flushing • unable to sweat • feeling sick or vomiting • dry mouth • dry and/or itchy skin • constipation • difficulty urinating • impotence • reduced milk supply in breastfeeding women • blurred vision or other problems with your eyesight • headache • nervousness, drowsiness • dizziness • unable to sleep 	<p>Speak to your doctor or nurse if you have any of these less serious side effects and they worry you.</p> <p>These side effects are usually mild.</p>

<ul style="list-style-type: none"> • over-excitement in children • confusion, especially in elderly people • fever • feeling bloated • loss or alteration of taste • injection site reactions like itchy skin, swelling, pain. These side effects are rare. 	
---	--

Contains no antimicrobial preservative.

Do not take this medicine if you are allergic to any of these ingredients.

What ROBINUL injection looks like

ROBINUL injection comes in a clear glass 1 mL ampoule containing a clear, colourless solution. Available in packs of 5 single dose ampoules.

Australian Registration number: AUST R 163586

Serious side effects

Serious side effects	What to do
<ul style="list-style-type: none"> • rash, itching or hives on the skin, swelling of the face, lips, tongue or other parts of the body, shortness of breath, wheezing or trouble breathing • cold sweat, nausea, light headedness, discomfort in chest or other areas of the upper body like pain or discomfort in one or both arms, the back, neck, jaw or stomach. 	<p>Call your doctor immediately or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.</p> <p>Serious side effects are rare.</p>

Who distributes ROBINUL injection

Aspen Pharmacare Australia Pty Ltd
34-36 Chandos St
St Leonards NSW 2065
Australia

This leaflet was revised in October 2025.

Tell your doctor, nurse or pharmacist if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people.

Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at www.tga.gov.au/reporting-problems. By reporting side effects, you can help provide more information on the safety of this medicine.

Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.

7. Product details

This medicine is only available with a doctor's prescription.

What ROBINUL injection contains

Active ingredient (main ingredient)	<ul style="list-style-type: none"> • Each 1 mL ampoule contains 200 microgram/1 mL of glycopyrronium bromide.
Other ingredients (inactive ingredients)	<ul style="list-style-type: none"> • water for injections • sodium chloride • sodium hydroxide and/or hydrochloric acid (for pH adjustment).