

DUTASTERIDE/TAMSULOSIN-WGR 500/400

Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

1. Why am I taking this medicine?

DUTASTERIDE/TAMSULOSIN-WGR 500/400 contains the active ingredients dutasteride and tamsulosin hydrochloride. DUTASTERIDE/TAMSULOSIN-WGR 500/400 is used in men who have benign prostatic hyperplasia (BPH). For more information, see Section [1. Why am I taking this medicine?](#) in the full CMI.

2. What should I know before I take this medicine?

Do not use if you have ever had an allergic reaction to DUTASTERIDE/TAMSULOSIN-WGR 500/400, any other 5 alpha reductase inhibitors or any of the ingredients listed at the end of the CMI.

Talk to your doctor if you have any other medical conditions or take any other medicines. Talk to your doctor if your partner is pregnant or plans to become pregnant or is breastfeeding.

For more information, see Section [2. What should I know before I take this medicine?](#) in the full CMI.

3. What if I am taking other medicines?

Some medicines may interfere with DUTASTERIDE/TAMSULOSIN-WGR 500/400 and affect how it works.

A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

4. How do I take this medicine?

- The usual dosage is one capsule once per day.
- Take your medicine approximately 30 minutes after the same meal each day. Do not take DUTASTERIDE/TAMSULOSIN-WGR 500/400 on an empty stomach.

More instructions can be found in Section [4. How do I take this medicine?](#) in the full CMI.

5. What should I know while taking this medicine?

Things you should do	<ul style="list-style-type: none">• Remind any doctor, dentist or pharmacist you visit that you are taking DUTASTERIDE/TAMSULOSIN-WGR 500/400.• If you are having a blood test remind your doctor that you are taking DUTASTERIDE/TAMSULOSIN-WGR 500/400.• If you are having cataract surgery, tell your surgeon that you are taking DUTASTERIDE/TAMSULOSIN-WGR 500/400.
Things you should not do	<ul style="list-style-type: none">• Do not stop using this medicine suddenly or change the dose.• Do not donate blood whilst taking DUTASTERIDE/TAMSULOSIN-WGR 500/400 or for 6 months after stopping treatment.
Driving or using machines	<ul style="list-style-type: none">• Do not drive or operate machinery until you know how this medicine affects you.• This medicine can cause dizziness, light headedness and potential fainting due to a sudden drop in blood pressure upon standing up (orthostatic hypotension).
Looking after your medicine	<ul style="list-style-type: none">• Store your capsules in the bottle until it is time to take them.• Store the capsules below 25°C.

For more information, see Section [5. What should I know while taking this medicine?](#) in the full CMI.

6. Are there any side effects?

Side effects which have been reported include impotence (an inability to maintain an erection), decrease in libido (sex drive), difficulty with ejaculation, breast swelling or tenderness, difficulty in sleeping (insomnia), dizziness, constipation, diarrhoea, vomiting, a fast heart beat, feeling weak or a loss of strength, low blood pressure upon standing up, itchy, blocked or runny nose and allergic reactions.

For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

DUTASTERIDE/TAMSULOSIN-WGR 500/400

Active ingredient(s): *dutasteride 500 micrograms and tamsulosin hydrochloride 400 micrograms*

Consumer Medicine Information (CMI)

This leaflet provides important information about using DUTASTERIDE/TAMSULOSIN-WGR 500/400. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using DUTASTERIDE/TAMSULOSIN-WGR 500/400.**

Where to find information in this leaflet:

- [1. Why am I taking this medicine?](#)
- [2. What should I know before I take this medicine?](#)
- [3. What if I am taking other medicines?](#)
- [4. How do I take this medicine?](#)
- [5. What should I know while taking this medicine?](#)
- [6. Are there any side effects?](#)
- [7. Product details](#)

1. Why am I taking this medicine?

DUTASTERIDE/TAMSULOSIN-WGR 500/400 contains the active ingredients dutasteride and tamsulosin hydrochloride.

Dutasteride belongs to a group of medicines called 5-alpha reductase enzyme inhibitors. Tamsulosin hydrochloride belongs to a group of medicines called alpha-blockers.

DUTASTERIDE/TAMSULOSIN-WGR 500/400 is used to in men with benign prostatic hyperplasia (BPH).

BPH is a non-cancerous enlargement of the prostate gland which is located in the lower portion of the urinary bladder surrounding the urethra (urine carrying tube). In men with BPH, the prostate gland becomes large enough to squeeze the urine tube running through it. If this tube is squeezed it narrows, making it more difficult to pass urine and you may have some or all of the following symptoms:

- difficulty in starting to urinate
- an interrupted, weak urinary stream
- more frequent urination, especially at night
- feeling that you need to urinate right away
- leaking or dribbling
- a feeling that you cannot empty your bladder completely

As the disease progresses, untreated BPH can lead to an increased risk of complete blockage of urine flow (acute urinary retention) and/or the need for surgery.

Prostate growth is caused by a hormone in the blood called dihydrotestosterone (DHT). The dutasteride component of DUTASTERIDE/TAMSULOSIN-WGR 500/400 lowers DHT production in the body leading to a shrinkage of the enlarged prostate in most men. The tamsulosin hydrochloride component of DUTASTERIDE/TAMSULOSIN-

WGR 500/400 acts by relaxing the smooth muscle in your prostate gland making it easier to pass urine.

Just as your prostate becomes enlarged over a long period of time, reducing the size of your prostate and improving your symptoms may take time.

Your doctor may have prescribed this medicine for another reason. Ask your doctor if you are unsure why this medicine was prescribed for you.

This medicine is not addictive.

2. What should I know before I take this medicine?

Warnings

Do not use DUTASTERIDE/TAMSULOSIN-WGR 500/400 if:

- you are allergic to dutasteride, tamsulosin hydrochloride, or any of the ingredients listed at the end of this leaflet. Always check the ingredients to make sure you can use this medicine.
- you are allergic to any other 5 alpha reductase enzyme inhibitors
- you are a woman or child
- you have a history of a sudden drop in blood pressure upon standing up (known as orthostatic hypotension)
- you have liver or kidney disease
- you are taking other medications that relax the smooth muscle of your blood vessels (e.g. Flomaxtra, Minipress)
- the expiry date (EXP) printed on the pack has passed
- the packaging is torn or shows signs of tampering

Check with your doctor if you:

- have any other medical conditions
- suffer from any liver or kidney conditions
- are going to have cataract surgery
- are taking any medicines to lower your blood pressure (alpha blocker medicines that relax the smooth muscles within blood vessels)
- are taking an antifungal medicine (e.g. ketoconazole)
- you have an allergy to sulfur
- take any medicines for any other condition

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

Pregnancy and breastfeeding

Talk to your doctor if your partner is pregnant or intends to become pregnant or is breastfeeding.

Women who are pregnant or may be pregnant, and children, must avoid handling punctured or leaking capsules. If skin contact occurs wash the area immediately with soap and water.

If dutasteride is absorbed through the skin by a woman who is pregnant with a male baby, it may cause the baby to be born with abnormalities of the genital organs.

Dutasteride has been found in the semen of men taking dutasteride. If your partner is or may be pregnant you must avoid exposing her to your semen as dutasteride may affect the normal development of a male baby. **You must use a barrier form of contraception.**

3. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

Tell your doctor or pharmacist if you are allergic to any foods, dyes, preservatives or any other medicines.

Some medicines may interfere with DUTASTERIDE/TAMSULOSIN-WGR 500/400 and affect how it works.

- long-term combination of dutasteride with medicines that are potent inhibitors of the enzyme CYP3A4 (e.g. ritonavir, indinavir, nefazodone, itraconazole, ketoconazole) may increase the concentration of dutasteride in the blood
- antifungal medicines such as ketoconazole can lead to an increase in the concentration of tamsulosin hydrochloride in the blood
- antidepressants known as selective serotonin reuptake inhibitors (e.g. paroxetine) can lead to an increase in the concentration of tamsulosin hydrochloride in the blood

Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect DUTASTERIDE/TAMSULOSIN-WGR 500/400.

4. How do I take this medicine?

How much to take

- The usual dose is one capsule daily. Follow the instructions provided and use this medicine until your doctor tells you to stop.

When to take DUTASTERIDE/TAMSULOSIN-WGR 500/400

- DUTASTERIDE/TAMSULOSIN-WGR 500/400 should be taken approximately 30 minutes after the same meal each day. Do not take this medicine on an empty stomach.
- Swallow your capsule whole with some water.
- Do not crush or chew the capsules. The contents of the capsule may make your mouth or throat sore.

If you forget to take DUTASTERIDE/TAMSULOSIN-WGR 500/400

This medicine should be taken at the same time each day.

If you forget to take DUTASTERIDE/TAMSULOSIN-WGR 500/400 skip the dose you missed and take your next dose when you are meant to.

Do not take a double dose to make up for the dose you missed.

If you take too much DUTASTERIDE/TAMSULOSIN-WGR 500/400

If you think that you have used too much DUTASTERIDE/TAMSULOSIN-WGR 500/400, you may need urgent medical attention.

You should immediately:

- phone the Poisons Information Centre (**by calling 13 11 26**), or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

You should do this even if there are no signs of discomfort or poisoning.

5. What should I know while taking this medicine?

Things you should do

Tell your doctor if you:

- have not taken this medicine exactly as directed
- are having a blood test, tell your doctor you are taking DUTASTERIDE/TAMSULOSIN-WGR 500/400 as this medicine decreases your prostate-specific antigen (PSA) levels.

The PSA test helps your doctor to tell if you have prostate disease, including prostate cancer. If you have a higher than normal amount of PSA in your blood it could mean that you have a higher risk of prostate cancer. Men taking DUTASTERIDE/TAMSULOSIN-WGR 500/400 should have their PSA measured 6 months after starting treatment and then regularly after that. You could still be at risk for prostate cancer even though your PSA is lower. Your doctor can still use PSA to help detect prostate cancer, by comparing your test results each time you have a PSA test.

Remind any doctor, dentist or pharmacist you visit that you are taking this medicine.

Things to be careful of

If you are having a cataract operation and are taking or have previously taken this medicine, the pupil may dilate poorly and the iris (the coloured part of the eye) may become floppy during the operation (floppy iris syndrome). This can be managed if your surgeon knows that you are taking DUTASTERIDE/TAMSULOSIN-WGR 500/400 before you have the operation. If you are going to have eye surgery tell your surgeon that you are taking or have taken DUTASTERIDE/TAMSULOSIN-WGR 500/400.

In a clinical study of men at increased risk of developing prostate cancer, a serious form of prostate cancer was reported more often in men taking this medicine than men who did not take this medicine. The reasons for this is currently unknown but may be due to the design of the clinical study.

Be careful if you are switching from another tamsulosin containing medicine to DUTASTERIDE/TAMSULOSIN-WGR 500/400 as you may experience dizziness or light headedness when standing up.

If you feel dizzy or lightheaded at any time whilst taking this medicine, sit or lie down until the symptoms pass.

Things you should not do

- Do not stop using this medicine suddenly or change the dose.

Some men notice an early improvement in their symptoms however others may need to continue treatment for 3 to 6 months to see if DUTASTERIDE/TAMSULOSIN-WGR 500/400 works for them.

Continue to take this medicine as your symptoms improve.

If you do not take your medicine as directed it may interfere with your doctor's ability to monitor your PSA levels.

- Do not donate blood whilst taking this medicine or for 6 months after you stop treatment.
- Do not give this medicine to anyone else even if they have similar symptoms to you.
- Do not use this medicine to treat any other complaints.

Driving or using machines

Be careful before you drive or use any machines or tools until you know how DUTASTERIDE/TAMSULOSIN-WGR 500/400 affects you.

DUTASTERIDE/TAMSULOSIN-WGR 500/400 may cause some people to experience dizziness or light headedness.

Looking after your medicine

Follow the instructions on the bottle on how to take care of your medicine properly.

Store it in a cool dry place (below 25°C) away from moisture, heat or sunlight; for example, do not store it:

- in the bathroom or near a sink, or
- in the car or on window sills.

Store your capsules in the bottle until it is time to take them.

Keep it where young children cannot reach it.

Getting rid of any unwanted medicine

If you no longer need to use this medicine or it is out of date, take it to any pharmacy for safe disposal.

Do not use this medicine after the expiry date.

6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

In clinical studies where patients were taking both the individual components of DUTASTERIDE/TAMSULOSIN-WGR 500/400, the rate of heart failure (meaning your heart does not pump blood as well as it should) was higher than in patients taking the individual components alone. However, this was seen in less than 1% of patients.

If you are taking DUTASTERIDE/TAMSULOSIN-WGR 500/400, talk to your doctor about this and other possible side effects.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

Less serious side effects

Less serious side effects	What to do
<ul style="list-style-type: none"> impotence (inability to maintain an erection) decrease in libido (sex drive) difficulty with ejaculation or retrograde ejaculation (where the ejaculate goes back into the bladder instead of being squirted out) decrease in semen volume difficulty in sleeping (insomnia) dizziness constipation diarrhoea vomiting fast heart beat weakness or loss of strength hair loss (usually from the body) or hair growth fainting changes in mood, feeling depressed itchy, blocked or runny nose changes in your vision nose bleeds dry mouth 	<p>Speak to your doctor if you have any of these less serious side effects and they worry you.</p>

Serious side effects

Serious side effects	What to do
<ul style="list-style-type: none"> allergic reaction – signs include wheezing, difficulty breathing, swelling of the lips or mouth, a rash or hives on the skin and fainting breast swelling or tenderness. If this becomes troublesome or 	<p>Call your doctor straight away, or go straight to the Emergency Department at your nearest</p>

Serious side effects	What to do
<p>you notice breast lumps or nipple discharge talk to your doctor about these changes as they could be signs of a serious condition such as breast cancer</p> <ul style="list-style-type: none"> serious skin reactions with a widespread rash, blisters and peeling skin particularly around the mouth, nose, eyes and genitals (such as Stevens-Johnson syndrome) testicular pain and/or swelling low blood pressure upon standing (orthostatic hypotension). Take care when moving from a lying or sitting position to a standing position. prolonged erection unrelated to sexual activity 	<p>hospital if you notice any of these serious side effects.</p>

Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people.

Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at www.tga.gov.au/reporting-problems. By reporting side effects, you can help provide more information on the safety of this medicine.

Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.

7. Product details

This medicine is only available with a doctor's prescription.

What DUTASTERIDE/TAMSULOSIN-WGR 500/400 contains

Active ingredient (main ingredient)	<ul style="list-style-type: none"> dutasteride tamsulosin hydrochloride
Other ingredients (inactive ingredients)	<ul style="list-style-type: none"> propylene glycol monocaprylate, butylated hydroxytoluene, methacrylic acid – ethyl acrylate copolymer (1:1), sodium laurylsulfate, polysorbate 80, microcrystalline cellulose, dibutyl sebacate, silicon dioxide, calcium stearate,

	<ul style="list-style-type: none"> Capsugel Ink 10A2 Black ID: 109521, iron oxide black, iron oxide red, iron oxide yellow, titanium dioxide, gelatin, glycerol.
Potential allergens	Soya bean products

Do not take this medicine if you are allergic to any of these ingredients.

What DUTASTERIDE/TAMSULOSIN-WGR 500/400 looks like

DUTASTERIDE/TAMSULOSIN-WGR 500/400 capsules are an oblong, hard-shell gelatin capsules with a brown body and a beige cap imprinted with C001 in black ink.

DUTASTERIDE/TAMSULOSIN-WGR 500/400 is available in bottles of 30 capsules.

Not all pack sizes may be distributed in Australia.

AUST R 430737

Who distributes DUTASTERIDE/TAMSULOSIN-WGR 500/400

Sponsored by

Helix Pharmaceuticals Pty Ltd
C/-EGA Corporate Advisors Pty Ltd
Level 12, 468 St Kilda Road
Melbourne, VIC 3000
Australia

Distributed by

GM Pharma International Pty Ltd
Level 12, 468 St Kilda Road
Melbourne VIC 3004 Australia
Tel: 1800 936 140

This leaflet was prepared in October 2025.