

TENOFOVIR/EMTRICITABINE 300/200 APX

Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

This medicine is new or being used differently. Please report side effects. See the [full CMI](#) for further details. [Include if applicable]

1. Why am I using TENOFOVIR/EMTRICITABINE 300/200 APX?

TENOFOVIR/EMTRICITABINE 300/200 APX contains the active ingredient tenofovir disoproxil fumarate, also called tenofovir DF and emtricitabine or FTC. TENOFOVIR/EMTRICITABINE 300/200 APX is used to treat Human Immunodeficiency Virus-1 (HIV-1) infection in adults when taken in combination with other anti-HIV medicines.

For more information, see Section [1. Why am I using this medicine?](#) in the full CMI.

2. What should I know before I use TENOFOVIR/EMTRICITABINE 300/200 APX?

Do not use if you have ever had an allergic reaction to this medicine or any of the ingredients listed at the end of the CMI.

Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.

For more information, see Section [2. What should I know before I use this medicine?](#) in the full CMI.

3. What if I am taking other medicines?

Some medicines may interfere with TENOFOVIR/EMTRICITABINE 300/200 APX and affect how it works.

A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

4. How do I use TENOFOVIR/EMTRICITABINE 300/200 APX?

- Take this medicine exactly as prescribed. The usual dose is one tablet orally once daily.
- Take TENOFOVIR/EMTRICITABINE 300/200 APX at the same time each day to keep TENOFOVIR/EMTRICITABINE 300/200 APX blood levels constant.

More instructions can be found in Section [4. How do I use this medicine?](#) in the full CMI.

5. What should I know while using TENOFOVIR/EMTRICITABINE 300/200 APX?

Things you should do	<ul style="list-style-type: none">• Remind any doctor, dentist or pharmacist you visit that you are using TENOFOVIR/EMTRICITABINE 300/200 APX or if you are about to be started on any other medicines.• become pregnant or are trying to become pregnant.• If for any reason you have not taken your medicine exactly as prescribed.
Things you should not do	<ul style="list-style-type: none">• Do not stop using this medicine suddenly or change the dose without first checking with your doctor.• Do not give this medicine to anyone else even if their symptoms seem similar to yours
Driving or using machines	<ul style="list-style-type: none">• Be careful before you drive or use any machines or tools until you know how TENOFOVIR/EMTRICITABINE 300/200 APX affects you
Looking after your medicine	<ul style="list-style-type: none">• Keep TENOFOVIR/EMTRICITABINE 300/200 APX tablets in a cool, dry place where it stays below 30°C.• Discard any remaining product 45 days after first opening of the bottle.

TENOFOVIR/EMTRICITABINE 300/200 APX is not a cure for HIV infection or AIDS. While taking TENOFOVIR/EMTRICITABINE 300/200 APX you may still develop infections or other illnesses associated with HIV infection.

For more information, see Section [5. What should I know while using this medicine?](#) in the full CMI.

6. Are there any side effects?

All medicines can have side effects. Check with your doctor as soon as possible if you have any problems while taking TENOFOVIR/EMTRICITABINE 300/200 APX, even if you do not think the problems are connected with the medicine or are not listed in this leaflet.

For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

TENOFOVIR/EMTRICITABINE 300/200 APX

Active ingredient(s): *tenofovir disoproxil fumarate 300 mg emtricitabine 200 mg*

Consumer Medicine Information (CMI)

This leaflet provides important information about using TENOFOVIR/EMTRICITABINE 300/200 APX. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using TENOFOVIR/EMTRICITABINE 300/200 APX.**

Where to find information in this leaflet:

- [1. Why am I using this medicine?](#)
- [2. What should I know before I use this medicine?](#)
- [3. What if I am taking other medicines?](#)
- [4. How do I use this medicine?](#)
- [5. What should I know while using this medicine?](#)
- [6. Are there any side effects?](#)
- [7. Product details](#)

7. Why am I using this medicine?

TENOFOVIR/EMTRICITABINE 300/200 APX consists of two medicines combined in one tablet:

- tenofovir disoproxil fumarate, also called tenofovir DF
- emtricitabine or FTC

These are combined in one tablet to help control Human Immunodeficiency Virus (HIV) infection.

this medicine contains two active ingredients that belong to a group of antiviral medicines known as nucleoside and nucleotide reverse transcriptase inhibitors (NRTI).

TENOFOVIR/EMTRICITABINE 300/200 APX is used to treat Human Immunodeficiency Virus-1 (HIV-1) infection in adults when taken in combination with other anti-HIV medicines;

- to help reduce the risk of getting HIV infection when used with safer sex practices in:
- HIV-negative men who have sex with men, who are at high risk of getting infected with HIV-1 through sex.
- Male-female sex partners when one partner has HIV-1 infection and the other does not.

When TENOFOVIR/EMTRICITABINE 300/200 APX is used to treat HIV infection:

When used with other HIV-1 medicines to treat HIV-1 infection, TENOFOVIR/EMTRICITABINE 300/200 APX may help:

- Reduce the amount of HIV-1 in your blood. This is called “viral load”.

- Increase the number of CD4+ (T) cells in your blood that help fight off other infections.

Reducing the amount of HIV-1 and increasing the CD4+ (T) cells in your blood may help improve your immune system.

This may reduce your risk of death or infections that can happen when your immune system is weak.

Use in Children and Elderly

TENOFOVIR/EMTRICITABINE 300/200 APX is for adults.

Do not take TENOFOVIR/EMTRICITABINE 300/200 APX if you are”

- under the age of 18 years.
- over the age of 65 before discussing with your doctor.

Does TENOFOVIR/EMTRICITABINE 300/200 APX cure HIV OR AIDS?

TENOFOVIR/EMTRICITABINE 300/200 APX is not a cure for HIV infection or AIDS. While taking TENOFOVIR/EMTRICITABINE 300/200 APX you may still develop infections or other illnesses associated with HIV infection.

If you have HIV-1 infection, you must keep taking HIV-1 medicines to control HIV-1 infection and decrease HIV-related illnesses.

Does TENOFOVIR/EMTRICITABINE 300/200 APX reduce the risk of passing HIV to others?

TENOFOVIR/EMTRICITABINE 300/200 APX does not reduce the risk of passing HIV to others through sexual contact or blood contamination.

It is important to continue to take appropriate precautions to prevent passing HIV to others.

When TENOFOVIR/EMTRICITABINE 300/200 APX is used to reduce the risk of HIV infection

When used with safer sex practices, TENOFOVIR/EMTRICITABINE 300/200 APX may help to reduce the risk of getting HIV-1 infection if you are at high risk of getting HIV infection.

TENOFOVIR/EMTRICITABINE 300/200 APX reduces the risk of getting HIV-1 when you have been taking it before you are exposed to HIV-1.

For people using TENOFOVIR/EMTRICITABINE 300/200 APX to reduce the risk of getting HIV-1 infection:

TENOFOVIR/EMTRICITABINE 300/200 APX can only help reduce your risk of getting HIV-1 before you are infected.

Do not take TENOFOVIR/EMTRICITABINE 300/200 APX to help reduce your risk of getting HIV-1 if:

- you already have HIV-1 infection. If you are HIV-positive, you need to take other medicines with TENOFOVIR/EMTRICITABINE 300/200 APX to treat HIV. TENOFOVIR/EMTRICITABINE 300/200 APX by itself is not a complete treatment for HIV.
- you do not know your HIV-1 infection status. You may already be HIV-positive. You need to take other HIV-1 medicines with TENOFOVIR/EMTRICITABINE 300/200 APX to treat HIV-1.
- Many HIV-1 tests can miss HIV-1 infection in a person who has recently become infected. If you have flu-like symptoms, you could have recently become infected with HIV-1. Tell your healthcare provider if you had a flu-like illness within the last month before starting TENOFOVIR/EMTRICITABINE 300/200 APX or at any time while taking TENOFOVIR/EMTRICITABINE 300/200 APX. Symptoms of new HIV-1 infection include: tiredness, fever, joint or muscle aches, headache, sore throat, vomiting or diarrhoea, rash, night sweats or enlarged lymph nodes in the neck or groin.

8. What should I know before I use this medicine?

Warnings

Together with your doctor, you need to decide whether TENOFOVIR/EMTRICITABINE 300/200 APX is right for you.

Do not use TENOFOVIR/EMTRICITABINE 300/200 APX if:

- you are allergic to tenofovir, tenofovir DF, or any of the ingredients listed at the end of this leaflet.
- Always check the ingredients to make sure you can use this medicine.

Some of the symptoms of an allergic reaction may include:

- shortness of breath
- wheezing or difficulty breathing
- rash, itching or hives on the skin
- swelling of the face, lips, tongue or other parts of the body

The ingredients of this medicine are listed in the product description section of this leaflet.

Do not take this medicine if you are:

- *already taking any of the components of this medicine (tenofovir DF or emtricitabine). Do not take this medicine if you are taking lamivudine.*
- *taking adefovir dipivoxil.*
- *taking tenofovir alafenamide.*

Do not take this medicine after the expiry or “use by” date (EXP) printed on the bottle.

If you take it after the expiry date has passed, it may not work as well.

Do not take this medicine if the packaging is torn or shows signs of tampering.

If you are not sure whether you should be taking this medicine, talk to your doctor.

Check with your doctor if you:

- have any other medical conditions are allergic to foods, dyes, preservatives, or any other medicines.
- take any medicines for any other condition.
- have liver problems, including hepatitis B, or C virus infection.
- you are taking medication to treat your hepatitis C virus (HCV) infection (e.g. ledipasvir/sofosbuvir, sofosbuvir/velpatasvir)
- have kidney problems.
- have or have ever had abnormal bones or bone difficulties.

This medicine is only available from a pharmacist after it has been prescribed by a doctor who specializes in the treatment of HIV infection.

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

Pregnancy and breastfeeding

Check with your doctor if you are pregnant or intend to become pregnant.

Talk to your doctor if you are breastfeeding or intend to breastfeed.

The safe use of this medicine has not been demonstrated. For this reason, it is important that women of child-bearing age receiving treatment with this medicine use an effective method of contraception to avoid becoming pregnant.

If you are a female who is taking this medicine to reduce the risk of getting HIV-1 infection and you become pregnant while taking this medicine, talk to your healthcare provider to decide if you should keep taking this medicine.

The active substances in this medicine (tenofovir disoproxil fumarate and emtricitabine) have been found in breast milk at low concentrations.

Consequently, it is recommended that nursing mothers do not breast-feed during treatment with this medicine. In general, women infected with HIV should not breast-feed their infants in order to avoid transmission of HIV to their newborn infant.

If you wish to continue receiving treatment with this medicine, it is important you remain under the care of a hospital or doctor who specializes in the treatment of HIV infection.

- Avoid doing things that increase your risk of getting HIV-1 or spreading HIV-1 to other people:
- Do not have any kind of sex without protection. Always practice safer sex. Use latex or non-latex condoms, except lambskin, to reduce contact with semen, vaginal fluids, or blood.

- Do not share personal items that can have blood or body fluids on them, such as toothbrushes and razor blades.
- Do not share or re-use needles or other injection equipment.

Ask your healthcare provider if you have any questions about how to prevent getting HIV-1 or spreading HIV-1 to other people.

If you have a long-standing viral infection of your liver (hepatitis B) it may flare up when you stop taking this medicine.

This can cause serious illness particularly if your liver is already not working very well. If you have both HIV and hepatitis B, when you start taking this medicine and even after you stop, your doctor is likely to arrange tests from time to time to check how well your liver is working. This refers to any medical condition-specific, medicine-specific, and/or age-specific subheading(s) relevant for inclusion for certain categories/groups of users, as applicable to the medicine.

9. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins, or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

Tell your doctor if are taking:

- didanosine (also known as ddl).
- ledipasvir/sofosbuvir
- sofosbuvir/velpatasvir
- sofosbuvir/velpatasvir/voxilaprevir

Some medicines may interfere with TENOFOVIR/EMTRICITABINE 300/200 APX and affect how it works. Your doctor or pharmacist will be able to tell you what to do when taking TENOFOVIR/EMTRICITABINE 300/200 APX tablets with other medicines.

Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect TENOFOVIR/EMTRICITABINE 300/200 APX.

10. How do I use this medicine?

How much to take TENOFOVIR/EMTRICITABINE 300/200 APX

- Take one tablet once daily or as advised by your doctor.
- If you are not sure how much of this medicine you should take, check with your doctor or pharmacist. Do not change the amount of TENOFOVIR/EMTRICITABINE 300/200 APX you take unless told to do so by your doctor.
- Your doctor will tell you how much of this medicine to take and how often to take it. You will also find this

information on the label of your medicine container. Follow the instructions provided and use this medicine until your doctor tells you to stop.

Because your medicine helps to control your condition, but does not cure it, you will need to take this medicine every day. If you are taking TENOFOVIR/EMTRICITABINE 300/200 APX to reduce the risk of HIV-1 infection, take this medicine every day for the period of time as prescribed by your doctor.

- Do not miss any doses of TENOFOVIR/EMTRICITABINE 300/200 APX. Missing a dose lowers the amount of medicine in your blood.
- Do not stop taking TENOFOVIR/EMTRICITABINE 300/200 APX without first talking to your doctor.

When to take this medicine?

- This medicine is best taken with a meal or just afterwards, however taking it without food should not reduce the effectiveness of the medicine.

How to take TENOFOVIR/EMTRICITABINE 300/200 APX

If you have HIV infection your doctor will generally prescribe TENOFOVIR/EMTRICITABINE 300/200 APX in combination with other anti-HIV medicines.

Take this medicine exactly as prescribed. The usual dose is one tablet orally once daily. Take this medicine at the same time each day to keep TENOFOVIR/EMTRICITABINE 300/200 APX blood levels constant.

TENOFOVIR/EMTRICITABINE 300/200 APX is absorbed rapidly. Do not take another TENOFOVIR/EMTRICITABINE 300/200 APX dose if vomiting has occurred unless it occurs within 1 hour after taking.

If you forget to take this medicine:

- It is important to take the prescribed daily dose in order to get the maximum benefit of treatment.
- If you forget to take your daily dose of this medicine, take it as soon as you remember that day, and then go back to taking your medicine as you would normally.
- Do not take a double dose to make up for a forgotten dose.
- Do not take more than one TENOFOVIR/EMTRICITABINE 300/200 APX tablet in a day.

If you use too much (overdose)

If you think that you have used too much of this medicine, you may need urgent medical attention.

You should immediately:

- phone the Poisons Information Centre (**by calling 13 11 26**), or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

You should do this even if there are no signs of discomfort or poisoning.

11. What should I know while using TENOFOVIR/EMTRICITABINE 300/200 APX?

Things you should do

Call your doctor or pharmacist straight away if you:

- are taking TENOFOVIR/EMTRICITABINE 300/200 APX if you are about to be started on any other medicines.
- become pregnant or are trying to become pregnant.
- for any reason you have not taken your medicine exactly as prescribed.

If you are taking this medicine to reduce your risk of getting HIV

- Just taking this medicine may not keep you from getting HIV.
- You must continue using safer sex practices while you are taking this medicine to reduce your risk of getting HIV.
- You must stay HIV-negative to keep taking this medicine to reduce your risk of infection.
- Know your HIV status and the HIV status of your partners.
- Get tested for HIV at least every 3 months or when your healthcare provider tells you.
- Get tested for other sexually transmitted infections such as syphilis and gonorrhoea. These infections make it easier for HIV to infect you. This medicine will not stop you from getting these other infections.
- If you think you were exposed to HIV, tell your healthcare provider right away. They may want to do more tests to be sure you are still HIV-negative.
- Get information and support to help reduce risky sexual behavior.
- Have fewer sex partners.
- Do not miss any doses of this medicine. Missing doses may increase your risk of getting HIV infection.
- If you do become HIV-positive, you need more medicine than this medicine alone to treat HIV. This medicine by itself is not a complete treatment for HIV.
- If you have HIV and take only this medicine, over time your HIV may become harder to treat.

Remind any doctor, dentist or pharmacist or other health professionals you visit that you are using TENOFOVIR/EMTRICITABINE 300/200 APX.

Things you should not do:

- Do not stop using this medicine suddenly or change the dose without first checking with your doctor.
- Do not give this medicine to anyone else even if their symptoms seem similar to yours.
- Do not use TENOFOVIR/EMTRICITABINE 300/200 APX to treat any other complaints unless your doctor says to.

Driving or using machines

Be careful before you drive or use any machines or tools until you know how TENOFOVIR/EMTRICITABINE 300/200 APX affects you.

Looking after your medicine

- Keep TENOFOVIR/EMTRICITABINE 300/200 APX tablets in a cool, dry place where it stays below 30°C.
- Discard any remaining product 45 days after first opening of the bottle.
- Keep your TENOFOVIR/EMTRICITABINE 300/200 APX tablets in the bottle with the cap tightly closed until you take them.
- If you take TENOFOVIR/EMTRICITABINE 300/200 APX tablets out of their pack, they may not keep well.

Follow the instructions in the carton on how to take care of your medicine properly.

Store it in a cool dry place away from moisture, heat or sunlight; for example, do not store it:

- in the bathroom or near a sink, or
- in the car or on windowsills.
- Heat and dampness can destroy some medicines.

Keep it where young children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store them.

When to discard your medicine (as relevant)

If your doctor tells you to stop taking TENOFOVIR/EMTRICITABINE 300/200 APX, or the tablets have passed their expiry date, ask your pharmacist what to do with any tablets left over.

Getting rid of any unwanted medicine

If you no longer need to use this medicine or it is out of date, take it to any pharmacy for safe disposal.

Do not use this medicine after the expiry date.

12. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

Check with your doctor as soon as possible if you have any problems while taking TENOFOVIR/EMTRICITABINE 300/200 APX, even if you do not think the problems are connected with the medicine or are not listed in this leaflet.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

Less serious side effects

Less serious side effects	What to do
<ul style="list-style-type: none"> • diarrhoea • nausea • tiredness • headache • dizziness • depression • problems sleeping • abnormal dreams • rash • stomach-area (abdomen) pain • headache • decreased weight 	<p>Speak to your doctor if you have any of these less serious side effects and they worry you.</p>

Very Serious side effects

Very Serious side effects	What to do
<p>Allergy: Some people are allergic to medicines. If you have any of the following symptoms soon after taking your medicine, DO NOT TAKE ANY MORE:</p> <ul style="list-style-type: none"> • Skin troubles such as lumpy skin rash or “hives” • Swelling of the face, lips, mouth or throat which may cause difficulty in swallowing or breathing • Wheezing, chest pain or tightness • Fainting <p>Pancreatitis: These side effects may be due to a condition called pancreatitis which sometimes occurs in patients taking anti-HIV medicines:</p> <ul style="list-style-type: none"> • Severe stomach pain or cramps • Nausea • Vomiting <p>Serious Liver Problems (hepatotoxicity): These side effects may be due to a condition called hepatotoxicity with liver enlargement and fat deposits in the liver (steatosis) which sometimes occurs in patients taking anti-HIV medicines.</p> <ul style="list-style-type: none"> • Your skin or the white part of your eyes turns yellow (jaundice) • Your urine turns dark 	<p>Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.</p>

Very Serious side effects	What to do
<ul style="list-style-type: none"> • Your bowel movements (stools) turn light in colour • you don't feel like eating food for several days or longer • Nausea • Stomach-area pains <p>Lactic Acidosis These side effects may be due to a condition called lactic acidosis (build-up of an acid in the blood).</p> <p><i>Lactic acidosis can be a medical emergency and may need to be treated in the hospital.</i></p> <p>You may be more likely to get lactic acidosis or liver problems if you are female, very overweight (obese), or have been taking similar nucleoside analog-containing medicines, like TENOFOVIR/EMTRICITABINE 300/200 APX, for a long time.</p> <ul style="list-style-type: none"> • You feel very weak or tired • You have unusual (not normal) muscle pain • You have trouble breathing • You have stomach pain with nausea and vomiting • You feel cold, especially in your arms and legs • You feel dizzy or light headed • You have a fast or irregular heartbeat <p>Hepatic Flares</p> <ul style="list-style-type: none"> • Your doctor should test you to see if you have chronic hepatitis B infection before you start taking this medicine. • If you have chronic hepatitis B infection you should not stop your treatment without first discussing this with your doctor, as some patients have had blood tests or symptoms indicating a worsening of their hepatitis (“hepatic flare”) after stopping individual components (tenofovir disoproxil fumarate and emtricitabine) of TENOFOVIR/EMTRICITABINE 300/200 APX. • You may require medical exams and blood tests for several months after stopping treatment. 	<p>Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.</p>

Very Serious side effects	What to do
TENOFOVIR/EMTRICITABINE 300/200 APX is not approved for the treatment of hepatitis B, so you must discuss your hepatitis B therapy with your healthcare provider.	Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.

Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people. This list of side effects is not complete.

TENOFOVIR/EMTRICITABINE 300/200 APX may cause other serious side effects. Tell your doctor if you notice anything else that is making you feel unwell, even if it is not on this list.

New and worse kidney problems

If you have had kidney problems in the past or need to take another drug that can cause kidney problems, your healthcare provider may need to perform additional blood tests to check your kidneys.

Bone problems

Bone problems can happen in some people who take tenofovir disoproxil fumarate/emtricitabine. Bone problems include bone pain, or softening or thinning of bones, which may lead to fractures. Your healthcare provider may need to do tests to check your bones.

Signs and symptoms of inflammation

In some patients with advanced HIV infection (AIDS), signs and symptoms of inflammation from previous infections may occur soon after anti-HIV treatment is started. It is believed that these symptoms are due to an improvement in the body's immune response, which lets the body fight infections that may have been present with no obvious symptoms. If you notice any symptoms of infection, please tell your doctor immediately.

Some people may get other side effects while taking TENOFOVIR/EMTRICITABINE 300/200 APX. If you are concerned, talk to your doctor or pharmacist.

Ask your doctor or pharmacist if you don't understand anything in this list.

Do not be alarmed by this list of possible side-effects. Most of them are very rare and you may not experience any of them.

Reporting side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet.

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at

www.tga.gov.au/reporting-problems. By reporting side effects, you can help provide more information on the safety of this medicine.

Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.

13. Product details

This medicine is only available with a doctor's prescription.

What TENOFOVIR/EMTRICITABINE 300/200 APX contains

Active ingredient (main ingredient)	<ul style="list-style-type: none"> 300 mg tenofovir disoproxil fumarate equivalent to tenofovir disoproxil 245 mg 200 mg emtricitabine
Other ingredients (inactive ingredients)	<ul style="list-style-type: none"> croscarmellose sodium lactose monohydrate magnesium stearate microcrystalline cellulose pregelatinised starch <p>Film-coating:</p> <ul style="list-style-type: none"> hypromellose indigo carmine aluminium lake lactose monohydrate titanium dioxide triacetin
Potential allergens	Lactose Monohydrate

Do not take this medicine if you are allergic to any of these ingredients.

What TENOFOVIR/EMTRICITABINE 300/200 APX looks like

TENOFOVIR/EMTRICITABINE 300/200 APX tablets are blue coloured capsule shaped, biconvex film coated tablet, plain on both sides.

TENOFOVIR/EMTRICITABINE 300/200 APX tablets are supplied in bottles containing 30 tablets.

SPONSOR

Arrotex Pharmaceuticals Pty Ltd
15-17 Chapel Street
Cremorne VIC 3121
Australia

www.arrotex.com.au

Australian Registration Number: AUST R 327896

Date of preparation

This leaflet was prepared in January 2026.

