

OLANZAPINE RBX

Consumer Medicine Information (CMI) Summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

1. Why am I using Olanzapine RBX

Olanzapine RBX contains the active ingredient Olanzapine. It helps correct chemical imbalances in the brain, which may cause mental illness.

For more information, see Section [1. Why am I using Olanzapine RBX?](#) in the full CMI.

2. What should I know before I use Olanzapine RBX ?

Do not use if you have ever had an allergic reaction to Olanzapine or any of the ingredients listed at the end of the CMI.

Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.

For more information, see Section [2. What should I know before I use Olanzapine RBX?](#) in the full CMI.

3. What if I am taking other medicines?

Some medicines may interfere with Olanzapine RBX and affect how it works.

A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

4. How do I use Olanzapine RBX?

Your doctor will tell you how many tablets you should take. The dose your doctor will prescribe for you will usually be in the range 5 mg to 20 mg per day.

More instructions can be found in Section [4. How do I use Olanzapine RBX?](#) in the full CMI.

5. What should I know while using Olanzapine RBX ?

Things you should do	<ul style="list-style-type: none">• Remind any doctor, dentist or pharmacist you visit that you are using Olanzapine RBX.• Tell your doctor if you become pregnant while taking Olanzapine RBX.• Keep all of your doctor's appointments so that your progress can be checked.
Things you should not do	<ul style="list-style-type: none">• Do not stop taking your medicine, or change the dosage, without first checking with your doctor.• Do not give this medicine to anyone else, even if they have the same condition as you
Driving or using machines	<ul style="list-style-type: none">• Be careful before you drive or use any machines or tools until you know how Olanzapine RBX affects you. Olanzapine RBX may cause drowsiness in some people.
Drinking alcohol	<ul style="list-style-type: none">• Tell your doctor if you drink alcohol. The effects of alcohol could be made worse while taking Olanzapine RBX.
Looking after your medicine	<ul style="list-style-type: none">• Keep your medicine in a cool dry place where the temperature stays below 25°C.• Keep this medicine where children cannot reach it.

For more information, see Section [5. What should I know while using Olanzapine RBX??](#) in the full CMI.

6. Are there any side effects?

Like other medicines, Olanzapine RBX may cause some unwanted side effects. These are likely to vary from patient to patient, although not everybody gets them. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need urgent medical attention.

Tell your doctor if you notice anything unusual or if you are concerned about any aspect of your health, even if you think the problems are not connected with this medicine and are not referred to in this leaflet.

For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

OLANZAPINE RBX

Active ingredient(s): Olanzapine

Consumer Medicine Information (CMI)

This leaflet provides important information about using Olanzapine RBX. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using Olanzapine RBX.**

Where to find information in this leaflet:

- [1. Why am I using Olanzapine RBX?](#)
- [2. What should I know before I use Olanzapine RBX?](#)
- [3. What if I am taking other medicines?](#)
- [4. How do I use Olanzapine RBX?](#)
- [5. What should I know while using Olanzapine RBX ?](#)
- [6. Are there any side effects?](#)
- [7. Product details](#)

1. Why am I using Olanzapine RBX?

Olanzapine RBX contains the active ingredient Olanzapine. Olanzapine belongs to a group of medicines called antipsychotics. It helps to correct chemical imbalances in the brain, which may cause mental illness.

Olanzapine RBX is used to treat symptoms of schizophrenia and related psychoses. Schizophrenia is a mental illness with disturbances in thinking, feelings and behavior.

Olanzapine RBX alone, or in combination with lithium or valproate, is used for the short-term treatment of acute manic episodes associated with Bipolar I Disorder.

Olanzapine RBX is also a mood stabiliser that prevents further occurrences of the disabling high and low (depressed) extremes of mood associated with Bipolar I Disorder.

Bipolar I Disorder is a mental illness with symptoms such as feeling "high", having excessive amounts of energy, needing much less sleep than usual, talking very quickly with racing ideas and sometimes severe irritability.

Your doctor may have prescribed Olanzapine RBX for another reason.

Ask your doctor if you have any questions about why Olanzapine RBX has been prescribed for you.

This medicine is available only with a doctor's prescription.

Olanzapine RBX is not recommended for use in children under the age of 18 years as there is not enough information on its effects in this age group.

2. What should I know before I take Olanzapine RBX?

Warnings

Do not take Olanzapine RBX if:

- you are allergic to Olanzapine, or any of the ingredients listed at the end of this leaflet.
- Always check the ingredients to make sure you can use this medicine.
Signs of an allergic reaction may include a skin rash, itching, shortness of breath or swelling of the face, lips or tongue.
- the packaging is torn or shows signs of tampering or the tablets or wafers do not look quite right.
- the expiry date on the pack has passed. If you take this medicine after the expiry date has passed it may not work as well.

If you are not sure whether you should start taking Olanzapine RBX, talk to your doctor or pharmacist

Check with your doctor if you:

- **have had an allergic reaction to any medicine which you have taken previously to treat your current condition.**
- **you have or have had any medical conditions, especially the following:**
 - tumour of the pituitary gland (a small gland at the base of the brain)
 - disease of the blood or bone marrow with a reduced number of white or red blood cells
 - disease of the blood vessels of the brain, including stroke
 - prostate problems
 - kidney or liver disease
 - high blood sugar, diabetes or a family history of diabetes
 - lipid alterations such as high blood fat values (cholesterol and triglycerides)
 - breast cancer or a family history of breast cancer
 - paralytic ileus, a condition where the small bowel does not work properly
 - epilepsy (seizures or fits)
 - glaucoma, a condition in which there is usually a build up of fluid in the eye
 - heart disease, including irregular heart rhythm

- neuroleptic malignant syndrome, a reaction to some medicines with a sudden increase in body temperature, extremely high blood pressure and severe convulsions
- **DRESS**, a severe drug reaction to olanzapine, with a combination of three or more of the following: rash, eosinophilia (higher number of a group of white blood cells), fever, swollen lymph nodes or inflammation of liver, kidney, lung, or heart
- tardive dyskinesia, a reaction to some medicines with uncontrollable twitching or jerking movements of the arms and legs.
- Akathisia, a subjective feeling of restlessness and difficulty sitting still; sleep apnoea, a sleep disorder where a person has pauses in breathing or periods of shallow breathing during sleep.
- elderly with dementia- related psychosis
- previously had a stroke or transient ischaemic attack (mini- stroke) or if you have high blood pressure

Tell your doctor if

- you will be in a hot environment or do a lot of vigorous exercise. Olanzapine RBX may make you sweat less, causing your body to overheat.
- you smoke. Smoking may affect Olanzapine RBX or may affect how it works.
- you suffer from lactose intolerance (because Olanzapine RBX tablets contain lactose).

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

Pregnancy and breastfeeding

Check with your doctor if you are pregnant or intend to become pregnant.

Like most antipsychotic medicines, Olanzapine RBX is not recommended for use during pregnancy.

Newborn babies of mothers taking antipsychotic drugs (including Olanzapine RBX) during the last trimester of pregnancy are at risk of experiencing extrapyramidal neurological disturbances and/or withdrawal symptoms following delivery. These may include, but are not limited to agitation, tremor, muscle stiffness or weakness, drowsiness, feeding problems, and breathing difficulty. If there is a need to consider Olanzapine RBX during your pregnancy, your doctor will discuss with you the benefits and risks of using it.

Talk to your doctor if you are breastfeeding or intend to breastfeed.

It is recommended that you do not breast-feed while taking Olanzapine RBX.

3. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines may interfere with Olanzapine RBX and affect how it works. These include:

- medicines used to treat a fast or irregular heart beat (arrhythmia).
- medicines taken for anxiety or to help you sleep.
- medicines taken for depression
- carbamazepine, a medicine used for mood stabilisation and to treat epilepsy
- other centrally acting medicines (eg. Tranquillisers or strong painkillers)
- ciprofloxacin, a medicine used to treat bacterial infections
- medicines that lower blood pressure
- medicines used for Parkinson's disease
- medicines that can change the heart's electrical activity or make it more likely to change.
- activated charcoal

Your doctor or pharmacist has more information on medicines to be careful with or avoid while taking Olanzapine RBX.

Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect Olanzapine RBX.

4. How do I use Olanzapine RBX?

How much to take

Your doctor will tell you how many Olanzapine RBX tablets you should take. The dose your doctor will prescribe for you will usually be in the range 5 mg to 20 mg per day.

Your doctor may increase or decrease your dose in order to find the appropriate dose for your condition.

A lower starting dose may be prescribed for elderly patients over the age of 65 years.

When to take

Olanzapine RBX tablets should be taken once a day as advised by your doctor.

Take your prescribed dose at the same time each day.

Olanzapine RBX tablets can be taken with or without food.

How to take Olanzapine RBX

Olanzapine RBX tablets should be swallowed whole with a glass of water.

Follow the instructions provided and do not stop taking

Olanzapine RBX just because you feel better. It is important that you do NOT stop taking Olanzapine RBX unless your doctor tells you.

If you forget to use Olanzapine RBX

Olanzapine RBX should be used regularly at the same time each day. If you miss your dose at the usual time, and

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Otherwise, take it as soon as you remember then go back to taking your medicine as you would normally.

Do not take a double dose to make up for the dose you missed.

If you are not sure what to do, ask your doctor or pharmacist.

If you use too much Olanzapine RBX

If you have taken too much Olanzapine RBX, the most common signs are fast heart beat, agitation/aggression, difficulty speaking, uncontrollable movements and sleepiness (sedation).

If you think that you have used too much Olanzapine RBX, you may need urgent medical attention.

You should immediately:

- phone the Poisons Information Centre (by calling **13 11 26**), or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

You should do this even if there are no signs of discomfort or poisoning.

5. What should I know while using Olanzapine RBX?

Things you should do

It is important that you remember to take Olanzapine RBX daily and at the dose prescribed by your doctor.

Tell all doctors, dentists and pharmacists who are treating you that you are taking Olanzapine RBX.

If you are about to be started on any new medicine, remind your doctor, dentist or pharmacist that you are taking Olanzapine RBX.

Tell your doctor immediately if you become pregnant while taking Olanzapine RBX.

Keep all of your doctor's appointments so that your progress can be checked.

- Your doctor should monitor your weight while you are taking Olanzapine RBX.
- Patients with diabetes or who have a higher chance of developing diabetes should have their blood sugar checked often.
- Your doctor may request you have a blood test

from time to time to monitor your cholesterol levels.

- If you are over 65, your doctor may measure your blood pressure from time to time.

Tell your doctor if you are female and your monthly periods are absent for six months or more.

All thoughts of suicide or violence must be taken seriously. Talk to your doctor or mental health professional if you have thoughts about death, suicide, self-harm, or harming others.

These may be signs of changes or worsening in your mental illness.

Things you should not do

Do not stop taking Olanzapine RBX, or lower the dosage, even if you are feeling better, without checking with your doctor.

Do not give Olanzapine RBX to anyone else, even if their symptoms seem similar or they have the same condition as you. Your doctor has prescribed Olanzapine RBX for you and your condition

Things to be careful of

If Olanzapine RBX makes you feel light-headed, dizzy or faint, be careful when getting up from a sitting or lying position. Getting up slowly may help.

If outdoors, wear protective clothing and use at least a 30+ sunscreen.

Olanzapine RBX may cause your skin to be much more sensitive to sunlight than it is normally. Exposure to sunlight may cause a skin rash, itching, redness, or severe sunburn.

If your skin does appear to be burning, tell your doctor.

Make sure you keep cool in hot weather and keep warm in cool weather. Olanzapine RBX may affect the way your body reacts to temperature changes.

Antipsychotics have the potential to cause cardiac complications and sudden cardiac death.

Driving or using machines

Be careful before you drive or use any machines or tools until you know how Olanzapine RBX affects you.

Olanzapine RBX may cause drowsiness in some people.

Drinking alcohol

Tell your doctor if you drink alcohol.

The effects of alcohol could be made worse while taking Olanzapine RBX.

Your doctor may suggest you avoid alcohol while you are being treated with Olanzapine RBX.

Looking after your medicine

Keep your tablets in the blister pack until it is time to take them.

Follow the instructions in the carton on how to take care of your medicine properly.

Store it in a cool dry place where the temperature stays below 25°C, away from moisture, heat or sunlight; for example, do not store it:

- in the bathroom or near a sink, or
- in the car or on window sills.

Keep it where young children cannot reach it. A locked cupboard at least one and a half meters above the ground is a good place to store medicines.

Getting rid of any unwanted medicine

If you no longer need to use this medicine or it is out of date, take it to any pharmacy for safe disposal.

Do not use this medicine after the expiry date.

6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

Like other medicines, Olanzapine RBX may cause some unwanted side effects. These are likely to vary from patient to patient. Some side effects may be related to the dose of Olanzapine RBX. Accordingly, it is important that you tell your doctor as soon as possible about any unwanted effects. Your doctor may then decide to adjust the dose of Olanzapine RBX you are taking.

Some people may feel dizzy in the early stages of treatment, especially when getting up from a lying or sitting position. This side effect usually passes after taking Olanzapine RBX for a few days.

Less serious side effects

Less serious side effects	What to do
<ul style="list-style-type: none"> • drowsiness • unusual tiredness or weakness • fever • restlessness or difficulty sitting still • increased appetite, weight gain • constipation, bloating • dry mouth • swelling of your hands, feet and ankles 	<p>Speak to your doctor if you have any of these less serious side effects and they worry you.</p>

<ul style="list-style-type: none"> • aching joints • nose bleeds • dizziness, confusion, forgetfulness. • speech disorder • sleepwalking • sleep eating 	
---	--

Serious side effects

Uncommon side effects	What to do
<ul style="list-style-type: none"> • symptoms of sunburn (such as redness, itching, swelling or blistering of the skin) which occur more quickly than normal • rash or allergic reaction. • slow heart beat • changes in sexual functioning or sex drive in men or women. • prolonged and/or painful erection • unusual secretion of breast milk • breast enlargement in men or women. • symptoms of high sugar levels in the blood (including passing large amounts of urine, excessive thirst, having a dry mouth and skin and weakness). These may indicate the onset or worsening of diabetes. • lipid alterations, such as high blood fat values (cholesterol and triglycerides) • reaction following abrupt discontinuation (profuse sweating, nausea or vomiting) • absence of menstrual periods and changes in the regularity of menstrual periods • involuntary passing of urine or difficulty in initiating urination • unusual hair loss or thinning. 	<p>Tell your doctor as soon as possible if you notice any of these uncommon side effects. You may need medical attention.</p>

Very serious side effects

Very serious side effects	What to do
<ul style="list-style-type: none"> • Sudden signs of an allergic reaction such as a skin rash, itching, shortness of breath or swelling of the face, lips or tongue. • frequent infections such as fever, severe chills, sore throat or mouth ulcers • bleeding or bruising more easily than normal. • painful swollen leg, chest pain, or shortness of breath as these can be signs of blood clots in the lungs or legs • seizures, fits or convulsions • yellowing of the skin and/or eyes 	<p>Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of these very serious side effects. You may need urgent medical</p>

<ul style="list-style-type: none"> • nausea, vomiting, loss of appetite, generally feeling unwell, fever, itching • severe upper stomach pain often with nausea and vomiting (inflammation of the pancreas) • worm-like movements of the tongue, or other uncontrolled movements of the tongue, mouth, cheeks, or jaw which may progress to the arms and legs • sudden increase in body temperature, sweating, fast heart beat, muscle stiffness, high blood pressure and convulsions • sudden decrease in body temperature, intense trembling, lethargy, confusion, slow heartbeat and slow breathing • sharp chest pain, coughing of blood, or sudden shortness of breath • pain/tenderness in the calf muscle area • muscle pain, muscle weakness and brown urine • heart attack • heart palpitations and dizziness, which may lead to collapse. • fast breathing, shortness of breath, fever with chills, feeling tired or weak, chest pain while coughing, fast heartbeat. You may have pneumonia. • fever or swollen glands, especially if they occur together with or shortly after a skin rash. 	<p>attention or hospitalisation.</p>
---	---

The following additional side effects may occur in some groups of people taking Olanzapine RBX:

Elderly patients with dementia-related psychosis

- unusual manner of walking
- falls
- pneumonia
- involuntary passing of urine.
- Stroke
- transient ischemic attack – symptoms maybe, but not limited to, paralysis in face, arm or leg.

Parkinson's disease psychosis

Some patients with Parkinson's disease may hallucinate (see, feel or hear things that are not there) or develop worsening symptoms of Parkinson's disease.

In combination with lithium or valproate

Patients with bipolar mania taking Olanzapine RBX in combination with lithium or valproate may notice the following additional side effects:

- tremors
- speech disorder.

- dry mouth
- increased appetite
- weight gain

Tell your doctor if you notice anything unusual or if you are concerned about any aspect of your health, even if you think the problems are not connected with this medicine and are not referred to in this leaflet.

Some side effects, such as changes to liver function, blood cell counts, cholesterol or triglycerides can occur. These can only be found when your doctor does tests from time to time to check your progress.

Do not be alarmed by this list of side effects. Your doctor has weighed the risks of using this medicine against the benefits they expect it will have for you.

Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people.

Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at www.tga.gov.au/reporting-problems. By reporting side effects, you can help provide more information on the safety of this medicine.

Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.

7. Product details

This medicine is only available with a doctor's prescription.

What Olanzapine RBX contains

Active ingredient (main ingredient)	Olanzapine
Other ingredients (inactive ingredients)	magnesium stearate lactose hypromellose, microcrystalline cellulose, magnesium stearate
Potential allergens	Sugars as lactose

Do not take this medicine if you are allergic to any of these ingredients.

What Olanzapine RBX looks like

OLANZAPINE RBX 2.5 mg Tablets are light yellow to yellow, round, biconvex tablets, debossed with 'O4' on one side and plain on the other side. AUST R 157856

OLANZAPINE RBX 5 mg Tablets are light yellow to yellow, round, biconvex tablets, debossed with 'O5' on one side and plain on the other side. AUST R 157853

OLANZAPINE RBX 7.5 mg Tablets are light yellow to yellow, round, biconvex tablets, debossed with 'O6' on one side and plain on the other side. AUST R 157860

OLANZAPINE RBX 10 mg Tablets are light yellow to yellow, round, biconvex tablets, debossed with 'O7' on one side and plain on the other side. AUST R 157859

All strengths are available in blister packs containing 28 tablets.

Who distributes Olanzapine RBX

Sun Pharma ANZ Pty Ltd

12 Waterloo Road

Macquarie Park Sydney NSW 2113

Australia

customerservice.aus@sunpharma.com

Tel: 1800 726 229

This leaflet was prepared in Nov 2025