

RISPERIDONE INTAS

Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

1. Why am I using RISPERIDONE INTAS?

RISPERIDONE INTAS contains the active ingredient risperidone. Risperidone belongs to a group of medicines called antipsychotic agents, which improve the symptoms of certain types of mental illness.

For more information, see Section [1. Why am I using RISPERIDONE INTAS?](#) in the full CMI.

2. What should I know before I use RISPERIDONE INTAS?

Do not use if you have ever had an allergic reaction to risperidone or any of the ingredients listed at the end of the CMI.

Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.

For more information, see Section [2. What should I know before I use RISPERIDONE INTAS?](#) in the full CMI.

3. What if I am taking other medicines?

Some medicines may interfere with RISPERIDONE INTAS and affect how it works.

A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

4. How do I use RISPERIDONE INTAS?

- If you are about to be started on any new medicine, tell your doctor or pharmacist that you are taking risperidone.
- Tell any other doctors, dentists and pharmacists who are treating you that you are taking this medicine.

More instructions can be found in Section [4. How do I use RISPERIDONE INTAS?](#) in the full CMI.

5. What should I know while using RISPERIDONE INTAS?

Things you should do	<ul style="list-style-type: none">• Tell your doctor or pharmacist that you are taking risperidone.• If you become pregnant or start breastfeeding while you are taking risperidone, tell your doctor immediately.• If you are going to have surgery, tell the surgeon or anesthetist that you are using this medicine.
Things you should not do	<ul style="list-style-type: none">• Do not take your medicine to treat any other complaints unless your doctor tells you to.• Do not give this medicine to anyone else, even if they have the same condition as you.• Do not stop taking your medicine, or change the dosage, without first checking with your doctor. Do not use more than the recommended dose.
Driving or using machines	<ul style="list-style-type: none">• Be careful when driving or operating machinery until you know how this medicine affects you.• Risperidone may cause dizziness, drowsiness or light-headedness in some people, especially after the first dose.• Make sure you know how you react to risperidone before you drive a car, operate machinery, or do anything else that could be dangerous if you feel drowsy or dizzy.
Drinking alcohol	<ul style="list-style-type: none">• Do not drink alcohol whilst you are taking this medicine.• Risperidone can increase the effects of alcohol.
Looking after your medicine	<ul style="list-style-type: none">• Keep your tablets in a cool dry place where the temperature stays below 25°C.• Do not store your medicine, or any other medicine, in the bathroom or near a sink. Do not leave it on a window sill or in the car.

For more information, see Section [5. What should I know while using RISPERIDONE INTAS?](#) in the full CMI.

6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention. For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

RISPERIDONE INTAS

Active ingredient(s): *Risperidone*

Consumer Medicine Information (CMI)

This leaflet provides important information about using RISPERIDONE INTAS. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using RISPERIDONE INTAS.**

Where to find information in this leaflet:

- [1. Why am I using RISPERIDONE INTAS?](#)
- [2. What should I know before I use RISPERIDONE INTAS?](#)
- [3. What if I am taking other medicines?](#)
- [4. How do I use RISPERIDONE INTAS?](#)
- [5. What should I know while using RISPERIDONE INTAS?](#)
- [6. Are there any side effects?](#)
- [7. Product details](#)

1. Why am I using RISPERIDONE INTAS?

RISPERIDONE INTAS contains the active ingredient risperidone. Risperidone belongs to a group of medicines called antipsychotic agents, which improve the symptoms of certain types of mental illness.

Risperidone is used for:

- treatment of sudden (acute) and long-term (chronic) schizophrenia and other types of related psychoses. These are disorders related to thought, feeling and/or action
- short term treatment of acute mania associated with Bipolar 1 Disorder – symptoms of this condition may include elevated, expansive or irritable mood, inflated self-esteem, decreased need for sleep, pressured speech, racing thoughts, distractibility or poor judgment including disruptive or aggressive behaviours
- short term treatment of behavioural problems in patients with a decline in mental ability (dementia) – these problems include aggression through words or actions, morbid suspiciousness, agitation or wandering
- conduct and other disruptive disorders such as aggression, impulsiveness and self-injury in adults, adolescents and children aged over 5 years old who are intellectually disabled
- certain behaviours seen in children and adolescents with autism.

How it works

- Risperidone helps to correct a chemical imbalance in the brain associated with the above conditions.
- Ask your doctor if you have any questions about why this medicine has been prescribed for you.
- Your doctor may have prescribed this medicine for another reason.
- This medicine is available only with a doctor's prescription.

There is no evidence this medicine is addictive.

2. What should I know before I use RISPERIDONE INTAS?

Warnings

Do not use RISPERIDONE INTAS if:

- you are allergic to risperidone, or any of the ingredients listed at the end of this leaflet.

Some of the symptoms of an allergic reaction may include:

- shortness of breath
- wheezing or difficulty breathing
- swelling of the face, lips, tongue, throat or other parts of the body
- skin rash, itching or hives on the skin

Always check the ingredients to make sure you can use this medicine.

Do not use this medicine after the expiry date printed on the pack or the packaging is torn or shows signs of tampering.

If it has expired or is damaged, return it to your pharmacist for disposal.

If you are not sure whether you should start taking this medicine, talk to your doctor.

Check with your doctor if you:

- have allergies to any other medicines, foods, preservatives or dyes.
- take any medicines for any other condition
- have or have had any of the following medical conditions:
 - Irregular heart rhythm, abnormalities in electrical activity of the heart, high or low blood pressure, or you have had a heart attack or stroke in the past.
 - unusual excessive sweating or diarrhoea, dehydration or problems with your body temperature regulation
 - prone to dizziness when standing up from a lying or sitting position

- kidney or liver problems
- Parkinson's disease (a progressive movement and thinking disorder that tends to affect older people)
- dementia or Lewy Body dementia older people suffering dementia may be at increased risk of stroke or death with Risperidone
- sugar diabetes
- unusual thirst, tiredness, blurred vision, upset stomach or need to urinate – common signs of high blood sugars
- epilepsy or seizures or fits
- restlessness or difficulty sitting still
- continuous and/or painful erections (called 'priapism')
- intraoperative iris syndrome (a complication that may occur during cataract extraction)
- Involuntary movements or unusual restlessness or difficulty sitting still
- suicidal thoughts or past suicide attempts
- low volume of blood in your body (hypovolaemia) – seen by cold hands and feet, light headedness, infrequent urination, increased heart rate, and weakness
- low blood potassium levels (hypokalaemia)
- breast cancer
- cancer of the pituitary gland
- diabetes, high or low blood sugar (you may need to monitor your blood glucose levels closely)
- tardive dyskinesia (a reaction to some medicines with uncontrollable twitching or jerking movements of the tongue, face, mouth, jaw, arms and legs)
- neuroleptic malignant syndrome (a serious reaction to some medicines with a sudden increase in body temperature, very fast heartbeat, extremely high or low blood pressure and severe muscle stiffness or fits)
- blood clots, Tell your doctor if you or someone else in your family has a history of blood clots. Blood clots in the lungs and legs can occur with Risperidone. Blood clots in the lungs can result in death.
- low white blood cell count

Tell your doctor if you are planning to have surgery (especially of the eye), dental treatment or an anesthetic.

- If you are having eye surgery, your doctor will need to assess whether you are at risk of a surgical complication (called 'Intraoperative Floppy Iris Syndrome).
- Tell your doctor if you will be in a hot environment or do a lot of vigorous exercise.
- Risperidone may make you sweat less, causing your body to overheat.

If you have not told your doctor about any of the above, tell them before you take this medicine.

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and

how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

Pregnancy and breastfeeding

Do not take this medicine if you are pregnant.

Do not take risperidone whilst pregnant until you and your doctor have discussed the risks and benefits involved.

Newborn babies of mother taking risperidone in their last trimester may be at risk of having difficulty feeding or breathing, shaking, muscle stiffness and/or weakness, sleepiness or agitation.

Do not breastfeed if you are taking this medicine.

Risperidone passes into breast milk and there is a possibility that your baby may be affected.

Check with your doctor if you are pregnant or breast feeding or intend to become pregnant and breast feed.

Do not take risperidone until you and your doctor have discussed the risks and benefits involved.

3. WHAT IF I AM TAKING OTHER MEDICINES?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines may interact with risperidone. These include:

- sleeping tablets, tranquillisers, strong pain-killers or antihistamines which may make you drowsy, certain antidepressants and alcohol
- medicines to treat Parkinson's disease or a tremor
- medicines that increase the activity of the central nervous system (e.g. psychostimulants such as methylphenidate)
- medicines to treat epilepsy such as carbamazepine (also used to treat trigeminal neuralgia)
- medicines to treat depression, panic disorder, anxiety, obsessive-compulsive disorder or premenstrual dysphoric disorder, such as tricyclic antidepressants (TCAs) and selective serotonin reuptake inhibitors (SSRIs) e.g. fluoxetine and paroxetine
- other medicines to treat mental illness or psychotic conditions
- diuretics, used to treat high blood pressure or swelling caused by the build-up of too much fluid e.g. frusemide – this combination may increase the risk of side effects or death in elderly people
- rifampicin, an antibiotic
- Verapamil, a medicine used to treat high blood pressure and/or abnormal heart rhythm.
- medicines used to treat heart problems or blood pressure
- medicines to relieve severe nausea and vomiting
- protease inhibitors, used to treat HIV infection (e.g. tipranavir, ritonavir)
- cimetidine and ranitidine, used for treating stomach ulcers or excess acid

- medicines which have an effect on the heart called QT prolongation
- medicines which affect the liver so that more of an enzyme called CYP3A4 is produced
- itraconazole and ketoconazole, medicines used to treat fungal infections.

These medicines may be affected by risperidone or may affect how well it works. You may need different amounts of your medicines, or you may need to take different medicines.

Your doctor and pharmacist have more information on medicines to be careful with or avoid while taking this medicine.

Other medicines not listed above may also interact with risperidone.

4. How do I use RISPERIDONE INTAS?

Follow all directions given to you by your doctor or pharmacist carefully. They may differ to the information contained in this leaflet.

If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

Taking it for the first time

At the start of treatment, you may have a fall in blood pressure making you feel dizzy on standing up, or your heart may beat faster.

Standing up slowly, especially when you get up from bed or chairs, will help your body get used to the change in position and blood pressure.

These symptoms should go away after a few days. Tell your doctor if they continue or worry you.

How much to take

Your doctor or pharmacist will tell you how much of this medicine you should take. This will depend on your condition and whether you are taking any other medicines.

- Risperidone may be taken as a single dose, once a day, or it may be taken in divided doses twice a day (in the morning and in the evening).
- Follow the instructions provided and use RISPERIDONE INTAS until your doctor tells you to stop.
- It is very important that you take the correct amount of risperidone, but this will vary from person to person. Your doctor will adjust the number and strength of the tablets until the desired effect is obtained.

For schizophrenia and related psychoses in adults, adolescents and children aged over 15 years old:

- The usual starting dose is one 1 mg tablet taken twice a day. This will be gradually increased by your doctor to suit your needs.
- From then on, the dose can be taken once a day or twice a day according to your doctor's instructions.

- For long-term treatment, 4 to 6 mg per day is usually sufficient, but your doctor will determine the dose most suitable for you.

Important Note: Never take more tablets than your doctor tells you to take.

- The effects of high doses are not yet known. Please double check with your doctor if your doctor prescribes more than 5 mg twice a day.
- Risperidone cannot be recommended for use in children with schizophrenia under 15 years as there is little experience with the product in this group.

For schizophrenia and related psychoses in elderly patients:

- The usual starting dose is one 0.5 mg tablet taken twice a day.
- The dose may be increased by 0.5 mg twice daily to 1 to 2 mg twice a day.

Patients who have impaired kidney or liver function:

- The usual starting dose is one 0.5 mg tablet taken twice a day. The dose may be increased by 0.5 mg twice daily to 1 to 2 mg twice a day.

For acute mania in patients with Bipolar 1 Disorder:

- The usual starting dose is one 2 mg tablet once a day. This dose can be adjusted by dose increases of 1 mg when needed every 24 hours. Most people feel better with doses between 2 mg and 6 mg a day. Your doctor may also decide you should take another drug called a mood stabiliser as well as risperidone.

For behavioural problems in people with dementia:

- The usual starting dose is 0.25 mg (half a 0.5 mg tablet) taken twice daily. This may be gradually increased by your doctor to suit your needs.
- From then on, the dose can be taken once a day or twice a day according to your doctor's instructions. For long-term treatment, 1 mg daily is the usual dose, but your doctor will determine the dose most suitable for you.

For conduct and other disruptive disorders in adults, adolescents and children aged over 5 years old who are intellectually disabled:

For people who weigh 50 kg or more, the usual starting dose is one 0.5 mg tablet taken once a day. The dose may be increased by 0.5 mg once every two days, to the usual dose of 0.5 to 1.5 mg once a day.

For people who weigh less than 50 kg, the usual starting dose is 0.25 mg (half a 0.5 mg tablet) taken once a day. The dose may be increased by 0.25 mg once every two days, to the usual dose of 0.25 to 0.75 mg once a day.

Your doctor will advise you on how much risperidone you need.

Risperidone cannot be recommended for use in children under 5 years or age with disruptive behaviour disorders as there is little experience with the product in this group.

For behavioural disorders associated with autism in children and adolescents:

- For people weighing less than 20 kg the usual starting dose is 0.25 mg (half a 0.5 mg tablet) daily. On Day 4, this dose can be increased to 0.5 mg.
- For people weighing 20 kg or more the usual starting dose is 0.5 mg (0.5 mg tablet or half a 1 mg tablet) daily. On Day 4, this dose can be increased to 1 mg.
- After 14 days, the doctor should check that the tablets are working. The doctor will say whether a higher dose is needed if the tablets do not seem to be have the desired effect. Your doctor will advise you on how much risperidone you need.
- In clinical trials, people with autism weighing less than 20 kg did not need more than 1.5 mg/day. People weighing over 20 kg did not need more than 2.5 mg of risperidone a day, and people weighing over 45 kg did not need more than 3.5 mg a day.
- If you feel sleepy, then the doctor may split up your daily dose so that you take half of it in the morning and half in the evening.

How to take it

Swallow the tablets whole with a full glass of water.

When to take it

- Take your medicine at about the same time each day.
- Taking it at the same time each day will have the best effect. It will also help you remember when to take it.
- It does not matter if you take this medicine before or after food.

How long to take it

- Continue taking your medicine for as long as your doctor tells you.
- This medicine helps to control your condition but does not cure it. It is important to keep taking your medicine even if you feel well.

If you forget to use RISPERIDONE INTAS

RISPERIDONE INTAS should be used regularly at the same time.

If you miss your dose at the usual time and its almost time to take your next dose, skip the dose you missed and take the next dose when you are meant to.

Otherwise take it as soon as you remember and then go back to taking your medicine as you would normally.

Do not take a double dose to make up for missed doses.

This may increase the chance of you experiencing side effects.

If you forget to take risperidone for 5 days or more, tell your doctor before starting your medicine again.

If you have trouble remembering to take your medicine, ask your pharmacist for some hints to help you remember

If you use too much RISPERIDONE INTAS

If you think that you have used too much RISPERIDONE INTAS, you may need urgent medical attention.

You should immediately:

- phone the Poisons Information Centre (by calling **13 11 26**), or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

You should do this even if there are no signs of discomfort or poisoning.

You may need urgent medical attention.

Symptoms of an overdose may include drowsiness, sleepiness, excessive trembling, excessive muscle stiffness, increased heart rate, and very low blood pressure causing fainting or unconsciousness.

5. What should I know while using RISPERIDONE INTAS?

Things you should do

If you are about to be started on any new medicine, tell your doctor or pharmacist that you are taking risperidone.

Tell any other doctors, dentists and pharmacists who are treating you that you are taking this medicine.

If you become pregnant or start breastfeeding while you are taking risperidone, tell your doctor immediately.

If you are going to have surgery, tell the surgeon or anaesthetist that you are using this medicine.

Keep all of your doctor's appointments so that your progress can be checked.

If a child is taking risperidone, tell the doctor if they do not seem to be developing normally.

Tell your doctor immediately if you notice any unusual movements of the tongue, or other uncontrolled movements of the tongue, mouth, cheeks or jaw which may progress to the arms and legs.

Your doctor may occasionally do tests to make sure risperidone is working and to prevent side effects.

Call your doctor straight away if you:

- There are symptoms of a condition called tardive dyskinesia, which may develop in people taking antipsychotic medicines, including risperidone. This condition is more likely to occur during long term treatment with risperidone, especially in elderly women. In very rare cases, this may be permanent. However, if detected early, these symptoms are usually reversible.
- Pre-menopausal women should tell their doctor if they do not have a period for more than six months while taking risperidone.
- Be careful during strenuous exercise or exposure to extreme heat. Try to drink plenty of water.

Remind any doctor, dentist or pharmacist you visit that you are using RISPERIDONE INTAS.

Things you should not do

- Do not take your medicine to treat any other complaints unless your doctor tells you to.
- Do not give this medicine to anyone else, even if they have the same condition as you.
- Do not stop taking your medicine, or change the dosage, without first checking with your doctor. Do not use more than the recommended dose.

Driving or using machines

Be careful before you drive or use any machines or tools until you know how Risperidone affects you.

Risperidone may cause dizziness, drowsiness or light-headedness in some people, especially after the first dose. Make sure you know how you react to risperidone before you drive a car, operate machinery, or do anything else that could be dangerous if you feel drowsy or dizzy.

Avoid excessive eating and try to eat a moderate diet.

Risperidone can cause weight gain.

Drinking alcohol

Tell your doctor if you drink alcohol.

Do not drink alcohol whilst you are taking this medicine.

Risperidone can increase the effects of alcohol.

Looking after your medicine

Keep your tablets in the pack until it is time to take them.

- If you take the tablets out of the pack, they may not keep well.
- Follow the instructions in the carton on how to take care of your medicine properly.
- Keep your tablets in a cool dry place where the temperature stays below 25°C.
- Heat and dampness can destroy some medicines.

Store it in a cool dry place away from moisture, heat or sunlight; for example, do not store it:

- in the bathroom or near a sink, or
- in the car or on window sills.

Keep it where young children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

When to discard your medicine

If your doctor tells you to stop taking this medicine or the expiry date has passed, ask your pharmacist what to do with any medicine that is left over.

Getting rid of any unwanted medicine

If you no longer need to use this medicine or it is out of date, take it to any pharmacy for safe disposal.

Do not use this medicine after the expiry date.

6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

Your doctor will decide whether any change in your treatment is needed.

This medicine helps most people, but it may have unwanted side effects in a few people. All medicines can have side effects. Sometimes they are serious, most of the time they are not.

Do not be alarmed by this list of side effects. You may not experience any of them.

Ask your doctor or pharmacist to answer any questions you may have.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

Less serious side effects

Less serious side effects	What to do
<ul style="list-style-type: none">• Difficulty thinking, working or carrying out your usual daily activities because of:• headache• trembling, muscle weakness, unsteadiness on your feet, lack of coordination or slow, shuffling walk (symptoms of Parkinsonism).• Lack of energy, drowsiness or excessive sleeping during the day, sleeplessness or difficulty in concentrating• Difficulty speaking• Blurred vision• Fainting• Dizziness• Any problems with confusion or unsteadiness• Pains in parts of your body, e.g. in the neck, back, ear, hands or feet• falling• muscle joint, nerve or movement changes such as:	<p>Speak to your doctor if you have any of these less serious side effects and they worry you.</p>

Less serious side effect	What to do
<ul style="list-style-type: none"> • shaking or trembling • fatigue or weakness • muscle stiffness • restlessness in the legs or difficulty sitting still • uncontrolled muscle spasms, twitching, jerky or writhing movements • muscle aches or pain • joint swelling or pain • walking abnormally or with difficulty • abnormal posture, such as rigid body movements and persistent abnormal positions of the body • behavioural changes such as: • irritability or agitation • unusual anxiety or nervousness • other changes such as: • cold or flu-like symptoms e.g. cough, blocked or runny nose, sneezing, sore throat • feeling of tension or fullness in the nose, cheeks and behind your eyes, sometimes with a throbbing ache, fever, stuffy nose and loss of the sense of smell (signs of sinusitis) • tiredness, headaches, being short of breath when exercising, dizziness and looking pale (signs of decreased red blood cells) • fever, chills, shortness of breath, cough, phlegm and occasionally blood (signs of pneumonia) • nosebleeds • discharge with itching of the eyes and crusty eyelids • unexplained weight gain • unexplained increase or decrease in appetite • indigestion, stomach discomfort or pain, diarrhoea or constipation • nausea or vomiting • dry mouth or excessive thirst • drooling • difficulty swallowing • acne • dry skin • rash, red skin or itchy skin • thickening of the skin resulting in warts, corns, calluses • skin infection • swelling of any part of your body, e.g. hands, ankles or feet • inability to or feeling burning pain when passing urine 	<p>Speak to your doctor if you have any of these less serious side effects and they worry you.</p>

Less serious side effects	What to do
<ul style="list-style-type: none"> • some loss of bladder control • bedwetting • frequent daytime urination in children • sexual function disturbances – problems with ejaculation • breast abnormalities – breast discomfort or swelling or unusual secretion of breast milk • missed or irregular menstrual periods • dizziness on standing up, especially when getting up from a sitting or lying down position • shortness of breath • chest pain or discomfort • an increase of CPK (creatine phosphokinase) in your blood, an enzyme which is sometimes released with muscle breakdown. <p>These can only be detected by blood tests that your doctor may ask to be done.</p>	<p>Speak to your doctor if you have any of these less serious side effects and they worry you.</p>

Serious side effects

Serious side effects	What to do
<p><u>Signs of heart or blood pressure problems including:</u></p> <ul style="list-style-type: none"> • Fainting, blurry vision, light-headedness or dizziness particularly on standing that persists despite sitting or lying down again • Very fast heart rate, slowed heart rate, heart rhythm irregularities. <p><u>Signs of lung problems including:</u></p> <ul style="list-style-type: none"> • Sudden shortness of breath, trouble breathing, wheezing or gasping when you breathe, light-headedness or dizziness <p><u>Signs of high blood sugar or diabetes such as:</u></p> <ul style="list-style-type: none"> • Unusual thirst, tiredness, upset stomach or need to urinate more often than usual <p><u>Body temperature changes such as:</u></p> <ul style="list-style-type: none"> • Fever • Unexplained high body temperature, excessive sweating or rapid breathing • Severe muscle stiffness or fits • Involuntary movements of the tongue, face, mouth, jaw, arms, legs or trunk 	<p>Call your doctor straight away or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.</p>

Serious side effects	What to do
<ul style="list-style-type: none"> Severe or life-threatening rash with blisters and peeling skin that may start in and around the mouth, nose, eyes and genitals and spread to other areas of the body (Stevens-Johnson syndrome or toxic epidermal necrolysis) Rash, itching or hives on the skin; shortness of breath, wheezing or difficulty breathing; swelling of the face, lips, tongue or other parts of the body. Sudden weakness or numbness of the face, arms or legs, especially on one side, or instances of slurred speech (these are called mini strokes) 	<p>Call your doctor straight away or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.</p>

Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people. The above list includes very serious side effects. You may need urgent medical attention or hospitalisation.

Other side effects not listed above may occur in some patients.

Do not hesitate to report any other side effects to your doctor or pharmacist. Do not be alarmed by this list of possible side effects.

You may not experience any of them

Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at www.tga.gov.au/reporting-problems. By reporting side effects, you can help provide more information on the safety of this medicine.

Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.

7. Product details

This medicine is only available with a doctor's prescription.

What RISPERIDONE INTAS contains

Active ingredient (main ingredient)	Each tablet contains 0.5 mg, 1 mg, 2 mg, 3 mg or 4 mg of risperidone as the active ingredient
Other ingredients (inactive ingredients)	<ul style="list-style-type: none"> lactose monohydrate maize starch microcrystalline cellulose hypromellose magnesium stearate colloidal anhydrous silica sodium lauryl sulfate purified talc propylene glycol titanium dioxide (0.5 mg, 2 mg, 3 mg and 4 mg tablets) iron oxide red (0.5 mg tablets) sunset yellow FCF aluminium lake (2 mg tablets) quinoline yellow aluminium lake (3 mg and 4 mg tablets) indigo carmine aluminium lake (4 mg tablets)
Potential allergens	Lactose

This medicine is gluten-free, contains sugars as lactose, tartrazine-free and free of other azo dyes.

Do not take this medicine if you are allergic to any of these ingredients.

What RISPERIDONE INTAS looks like

RISPERIDONE INTAS 0.5 mg tablets:

Brick red coloured, round, biconvex, film coated tablets plain on both sides

Blister pack of 20 or 60 tablets. AUST R 304250.

RISPERIDONE INTAS 1 mg tablets:

White to off white capsule shaped, biconvex, film coated tablets plain on both sides.

Blister pack of 60 tablets. AUST R 304251.

RISPERIDONE INTAS 2 mg tablets:

Light orange coloured, capsule shaped, biconvex, film coated tablets plain on both sides.

Blister pack of 60 tablets. AUST R 304252.

RISPERIDONE INTAS 3 mg tablets:

Light yellow coloured, oval shaped, biconvex, film coated tablets with break line on one side and plain on other side.

Blister pack of 60 tablets. AUST R 304253.

RISPERIDONE INTAS 4 mg tablets:

Green coloured, capsule shaped, biconvex, film coated tablets plain on both sides.

Blister pack of 60 tablets. AUST R 304254.

*Not all strengths, pack types and/or pack sizes may be available.

Who distributes RISPERIDONE INTAS

Arrotex Pharmaceuticals Pty Ltd

15-17 Chapel Street

Cremorne VIC 3121

www.arrotex.com.au

This leaflet was prepared in February 2026.