

LEFLUNOMIDE-WGR

Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

1. Why am I using LEFLUNOMIDE-WGR?

LEFLUNOMIDE-WGR contains the active ingredient leflunomide. LEFLUNOMIDE-WGR is used to treat rheumatoid or psoriatic arthritis. For more information, see Section [1. Why am I using LEFLUNOMIDE-WGR?](#) in the full CMI.

2. What should I know before I use LEFLUNOMIDE-WGR?

Do not use if you have ever had an allergic reaction to leflunomide, teriflunomide or any of the ingredients listed at the end of the CMI. Do not use if you are pregnant or plan to become pregnant, are not using reliable birth control, or are breastfeeding.

There are a number of other circumstances in which a person must not use this medicine. Check if these apply to you before taking LEFLUNOMIDE-WGR (see the full CMI for more details).

Talk to your doctor if you have any other medical conditions, take any other medicines. For more information, see Section [2. What should I know before I use LEFLUNOMIDE-WGR?](#) in the full CMI.

3. What if I am taking other medicines?

Some medicines may interfere with LEFLUNOMIDE-WGR and affect how it works. A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

4. How do I use LEFLUNOMIDE-WGR?

- The standard dose for this medicine is 100 mg per day for the first 3 days, and after that one 10 mg or 20 mg tablet daily.
- Swallow the tablets whole with a full glass of water, at about the same time each day. You can take it with or without food.

More instructions can be found in Section [4. How do I use LEFLUNOMIDE-WGR?](#) in the full CMI.

5. What should I know while using LEFLUNOMIDE-WGR?

Things you should do	<ul style="list-style-type: none">• Remind any doctor, dentist or pharmacist you visit that you are using LEFLUNOMIDE-WGR.• Call your doctor straight away if: you become pregnant while you are taking this medicine; you have an infection/notice a fever or signs of an infection; your skin becomes itchy or yellow / the whites of your eyes become yellow / if you start to bleed or bruise easily; you develop new /worsening cough or trouble breathing.• Tell your doctor before stopping contraception.
Things you should not do	<ul style="list-style-type: none">• Do not take more than the recommended dose unless your doctor tells you to.• Do not stop taking LEFLUNOMIDE-WGR, or lower the dosage, without checking with your doctor.
Drinking alcohol	<ul style="list-style-type: none">• Tell your doctor if you drink alcohol.• It is recommended that you minimise your alcohol intake while taking LEFLUNOMIDE-WGR.
Looking after your medicine	<ul style="list-style-type: none">• Store it in a cool dry place below 30°C. Store in a dry place and keep the container tightly closed.

For more information, see Section [5. What should I know while using LEFLUNOMIDE-WGR?](#) in the full CMI.

6. Are there any side effects?

Common side effects include: diarrhoea; nausea and vomiting; abdominal pain; weight loss; rashes, itchy skin; hair loss; unusual tiredness or weakness. Serious side effects include: fever; severe upper stomach pain; severe skin rash or sores in your mouth; pale skin, fatigue or increased infections and bruising; cough or trouble breathing; blisters and bleeding in the lips, eyes, mouth, nose and genitals; skin ulcer. Very serious side effects include: allergic reaction symptoms.

For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

LEFLUNOMIDE-WGR

Active ingredient: *leflunomide*

Consumer Medicine Information (CMI)

This leaflet provides important information about using LEFLUNOMIDE-WGR. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using LEFLUNOMIDE-WGR.**

Where to find information in this leaflet:

- [1. Why am I using LEFLUNOMIDE-WGR?](#)
- [2. What should I know before I use LEFLUNOMIDE-WGR?](#)
- [3. What if I am taking other medicines?](#)
- [4. How do I use LEFLUNOMIDE-WGR?](#)
- [5. What should I know while using LEFLUNOMIDE-WGR?](#)
- [6. Are there any side effects?](#)
- [7. Product details](#)

1. Why am I using LEFLUNOMIDE-WGR?

LEFLUNOMIDE-WGR contains the active ingredient leflunomide. LEFLUNOMIDE-WGR is an immunosuppressive medicine, which means that it works by reducing the activity of the immune system.

LEFLUNOMIDE-WGR is used to treat rheumatoid or psoriatic arthritis.

LEFLUNOMIDE-WGR helps to slow down the process of joint damage and to relieve the symptoms of the disease, such as joint tenderness and swelling, pain and morning stiffness.

LEFLUNOMIDE-WGR works by selectively interfering with the ability of white blood cells called lymphocytes to produce the disease response that ultimately leads to pain, inflammation and joint damage

2. What should I know before I use LEFLUNOMIDE-WGR?

Warnings

Do not use LEFLUNOMIDE-WGR if:

- are allergic to leflunomide, teriflunomide or any of the ingredients listed at the end of this leaflet. Always check the ingredients to make sure you can use this medicine
- have any diseases which reduce your body's natural defenses such as bacterial or viral infections
- have any diseases of the blood
- have any serious skin disorders
- have liver disease

- have a condition called hypoproteinaemia (when you do not have enough protein in your blood)
- are pregnant or plan to become pregnant
- are not using reliable birth control
- are breastfeeding

Check with your doctor if you:

- have or have had any other medical conditions, especially:
 - a decrease in the number of white blood cells
 - liver problems
 - kidney problems
 - chronic infections
 - an illness which lowered your body's resistance to disease
 - tuberculosis
 - lung problems, such as interstitial lung disease (an inflammation of lung tissue) which is a serious and potentially fatal disease
 - take any medicines for any other condition
 - are allergic to any other substances such as foods, preservatives or dyes
 - plan to have surgery

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

Pregnancy and breastfeeding

You must not become pregnant while taking LEFLUNOMIDE-WGR and for a certain period of time after stopping LEFLUNOMIDE-WGR.

- Tell your doctor immediately if you think you could be pregnant while taking LEFLUNOMIDE-WGR.
- Tell your doctor if you intend to become pregnant or father a child.

LEFLUNOMIDE-WGR may increase the risk of birth defects. To reduce any risk to the developing baby, you will need to stop taking LEFLUNOMIDE-WGR and may need to undergo a wash-out procedure. Your doctor will discuss the wash-out procedure with you.

Women of childbearing potential must use reliable contraception while taking LEFLUNOMIDE-WGR.

Do not take it if you are breastfeeding or planning to breastfeed.

- Tell your doctor if you are breast feeding or planning to breastfeed.

LEFLUNOMIDE-WGR passes into breast milk and therefore there is a possibility that the breastfed baby may be affected.

Use in Children

Do not give LEFLUNOMIDE-WGR to a child or adolescent.

LEFLUNOMIDE-WGR is not approved for use in children or adolescents under 18 years old.

3. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines may interfere with LEFLUNOMIDE-WGR and affect how it works.

- warfarin, a medicine used to stop blood clots
- some medicines used for diabetes
- some medicines used to treat epilepsy
- some medicines used for tuberculosis (TB)
- some medicines used to lower cholesterol
- some medicines used to treat depression
- some medicines used to treat respiratory conditions
- some medicines used to treat cancer
- some antibiotics
- some medicines used to treat autoimmune disorders
- some types of oral contraceptives

Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect LEFLUNOMIDE-WGR.

4. How do I use LEFLUNOMIDE-WGR ?

How much to take

- Your doctor will tell you how many tablets you will need to take each day. This depends on your condition and whether or not you are taking any other medicines.
- The standard dose for this medicine is 100 mg per day for the first 3 days, and after that one 10 mg or 20 mg tablet daily.
- Follow the instructions provided and use LEFLUNOMIDE-WGR until your doctor tells you to stop. The medicine helps control your condition, but it does not cure it. Therefore, you must take it every day.

When to take LEFLUNOMIDE-WGR

- LEFLUNOMIDE-WGR should be taken at about the same time each day.
 - Taking your tablets at the same time each day will have the best effect. It will also help you to remember when to take the tablets.
- It does not matter if you take LEFLUNOMIDE-WGR before or after food.

How to take LEFLUNOMIDE-WGR

- Swallow the tablets whole with a full glass of water.

If you forget to use LEFLUNOMIDE-WGR

LEFLUNOMIDE-WGR should be used regularly at the same time each day.

If you miss your dose at the usual time and there is still a long time to go before your next dose take it as soon as you remember. Then go back to taking it as you would normally.

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Do not take a double dose to make up for the dose you missed.

If you use too much LEFLUNOMIDE-WGR

If you think that you have used too much LEFLUNOMIDE-WGR, you may need urgent medical attention.

You should immediately:

- phone the Poisons Information Centre (by calling Australia 13 11 26 or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

You should do this even if there are no signs of discomfort or poisoning.

5. What should I know while using LEFLUNOMIDE-WGR?

In certain situations, for example, if you experience a serious side effect, you change your medication or you want to fall pregnant, your doctor will ask you to take medication that will help your body get rid of LEFLUNOMIDE-WGR faster.

Things you should do

- Tell your doctor before stopping contraception. You must continue using appropriate reliable contraception (the 'Pill' or condoms) while you are taking LEFLUNOMIDE-WGR.
- Tell your doctor if you develop symptoms such as pins and needles or tingling in the hands or feet or numbness or weakness of the arms and legs.
- Tell your doctor if you need to have a vaccination during treatment with this medicine or for 6 months after stopping treatment.
- Tell your doctor if you develop skin ulcer (open sore) during treatment with LEFLUNOMIDE-WGR (see section [6. Are there any side effects?](#))

Call your doctor straight away if:

- you become pregnant while you are taking this medicine. Stop taking it and tell your doctor or pharmacist immediately.
- you have an infection or notice a fever or signs of an infection
- your skin becomes itchy or yellow, if the whites of your eyes become yellow, or if you start to bleed or bruise easily. Stop taking it and tell your doctor or pharmacist immediately.

- You may be developing a liver problem. Your doctor may need to take blood samples to monitor the health of your liver and blood cells while you are taking LEFLUNOMIDE-WGR.
- you develop new or worsening symptoms such as a cough or trouble breathing.
 - Inflammation of the lung tissue which can be fatal, has been reported in some patients.

Remind any doctor, dentist or pharmacist you visit that you are using LEFLUNOMIDE-WGR.

Things you should not do

- Do not take more than the recommended dose unless your doctor tells you to.
- Do not stop taking LEFLUNOMIDE-WGR, or lower the dosage, without checking with your doctor.
- Do not give this medicine to anyone else, even if they have the same condition as you.
- Do not use this medicine to treat any other complaints unless your doctor tells you to.

Drinking alcohol

Tell your doctor if you drink alcohol.

The effects of alcohol could be made worse while taking LEFLUNOMIDE-WGR. It is recommended that you minimise your alcohol intake while taking LEFLUNOMIDE-WGR.

Looking after your medicine

Follow the instructions in the carton on how to take care of your medicine properly.

Store it in a cool dry place below 25°C away from moisture, heat or sunlight; for example, do not store it:

- in the bathroom or near a sink, or
- in the car or on window sills.

Keep your tablets in the container until it is time to take them. If you take the tablets out of the box or the blister pack they may not keep well.

Keep it where young children cannot reach it.

Getting rid of any unwanted medicine

If you no longer need to use this medicine or it is out of date, take it to any pharmacy for safe disposal.

Do not use this medicine after the expiry date.

6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

Less serious side effects

Less serious side effects	What to do
<p>Gut-related:</p> <ul style="list-style-type: none"> • diarrhoea • nausea and vomiting • abdominal pain • weight loss <p>Skin-related:</p> <ul style="list-style-type: none"> • paresthesia (numbness, tingling of the skin) • rashes, itchy skin <p>Other:</p> <ul style="list-style-type: none"> • hair loss • unusual tiredness or weakness • dizziness • headache • anorexia 	<p>Speak to your doctor if you have any of these less serious side effects and they worry you.</p>

Serious side effects

Serious side effects	What to do
<p>Skin-related:</p> <ul style="list-style-type: none"> • severe skin rash or sores in your mouth • blisters and bleeding in the lips, eyes, mouth, nose and genitals • pale skin, prone to infection or bruising and feeling tired • skin ulcer (a round, open sore in the skin through which the underlying tissues can be seen) <p>Other:</p> <ul style="list-style-type: none"> • signs and symptoms of severe infection (e.g. fever) • severe upper stomach pain, often with nausea and vomiting • new or worsening symptoms such as cough or trouble breathing, with or without fever 	<p>Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.</p>

Very serious side effects

Very serious side effects	What to do
<p>Allergic reaction-related:</p> <ul style="list-style-type: none"> • swelling of the face, lips, mouth or throat, which may cause difficulty in swallowing or breathing • hives • fainting • yellowing of the skin and eyes (jaundice) • thickened patches of red skin 	<p>Stop taking this medicine and tell your doctor immediately, or go to Accident and Emergency at your nearest hospital.</p>

Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people.

Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects online to the Therapeutic Goods Administration (Australia) at www.tga.gov.au/reporting-problems

By reporting side effects, you can help provide more information on the safety of this medicine.

Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.

7. Product details

This medicine is only available with a doctor's prescription.

What LEFLUNOMIDE-WGR contains

Active ingredient (main ingredient)	leflunomide
Other ingredients (inactive ingredients)	<ul style="list-style-type: none">• microcrystalline cellulose• lactose monohydrate• maize starch• povidone• crospovidone• colloidal anhydrous silica• magnesium stearate• Opadry II complete film coating system OY-LS-28908 White (10 mg only)• Opadry Aqueous Film Coating OY-SR-6497 Yellow (20 mg only)
Potential allergens	Contains sugars as lactose

Do not take this medicine if you are allergic to any of these ingredients.

LEFLUNOMIDE-WGR does not contain, sucrose, gluten, tartrazine or any other azo dyes.

What LEFLUNOMIDE-WGR looks like

LEFLUNOMIDE-WGR 10 mg tablets are white, round biconvex tablets.

LEFLUNOMIDE-WGR 10 mg tablets are supplied in bottles in packs of 30 tablets. AUST R 431390.

LEFLUNOMIDE-WGR 20 mg tablets are Yellow, round biconvex tablets with a scoreline on one side.

LEFLUNOMIDE-WGR 20 mg tablets are supplied in bottles in packs of 30 tablets. AUST R 431391.

Who distributes LEFLUNOMIDE-WGR

Wagner Pharmaceuticals Pty Ltd

6 Albert Street

Preston VIC 3072

Tel: 1800 936 140

This leaflet was prepared in November 2024