

APO-IRBESARTAN HCTZ

Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

1. Why am I taking APO-IRBESARTAN HCTZ?

APO-IRBESARTAN HCTZ contains the active ingredients irbesartan and hydrochlorothiazide. APO-IRBESARTAN HCTZ is used to treat high blood pressure. For more information, see Section [1. Why am I taking APO-IRBESARTAN HCTZ?](#) in the full CMI.

2. What should I know before I take APO-IRBESARTAN HCTZ?

Do not take if you have ever had an allergic reaction to APO-IRBESARTAN HCTZ or any of the ingredients listed at the end of the CMI.

Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.

For more information, see Section [2. What should I know before I take APO-IRBESARTAN HCTZ?](#) in the full CMI.

3. What if I am taking other medicines?

Some medicines may interfere with APO-IRBESARTAN HCTZ and affect how it works.

A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

4. How do I take APO-IRBESARTAN HCTZ?

- Your doctor will tell you how many tablets to take each day. The usual dose is one tablet per day.
- Swallow the tablet whole with a glass of water and take APO-IRBESARTAN HCTZ at about the same time each day.

More instructions can be found in Section [4. How do I take APO-IRBESARTAN HCTZ?](#) in the full CMI.

5. What should I know while taking APO-IRBESARTAN HCTZ?

Things you should do	<ul style="list-style-type: none">• Remind any doctor, dentist or pharmacist you visit that you are taking APO-IRBESARTAN HCTZ.• If you become pregnant while taking APO-IRBESARTAN HCTZ tell your doctor immediately.• Before being started on any new medicine, tell your doctor and pharmacist you are taking APO-IRBESARTAN HCTZ.• If you plan to have surgery that needs a general anaesthetic, tell your doctor or dentist that you are taking APO-IRBESARTAN HCTZ.• Make sure you drink enough water during exercise and hot weather when you are taking APO-IRBESARTAN HCTZ.• If you have excessive vomiting and/or diarrhoea while taking APO-IRBESARTAN HCTZ, tell your doctor.• If you feel light-headed or dizzy after your dose is increased, tell your doctor immediately.• Tell your doctor if you experience increased skin sensitivity.
Things you should not do	<ul style="list-style-type: none">• Do not stop taking APO-IRBESARTAN HCTZ or lower the dosage, without checking with your doctor.
Driving or using machines	<ul style="list-style-type: none">• Be careful driving or operating machinery until you know how APO-IRBESARTAN HCTZ affects you. APO-IRBESARTAN HCTZ may cause dizziness or light-headedness in some people. If this occurs do not drive.
Drinking alcohol	<ul style="list-style-type: none">• If you drink alcohol, dizziness or light-headedness may be worse.
Looking after your medicine	<ul style="list-style-type: none">• Keep APO-IRBESARTAN HCTZ tablets in a cool dry place where the temperature stays below 25°C.• Keep your tablets in the packaging until it is time to take them.

For more information, see Section [5. What should I know while taking APO-IRBESARTAN HCTZ?](#) in the full CMI.

6. Are there any side effects?

Common side effects: headache, dizziness or light-headedness (vertigo), unusual tiredness, weakness or fatigue. Shortness of breath when exercising or looking pale. Pain in the stomach or gut. Nausea and/or vomiting, sexual problems, low blood glucose levels. Refer to the full CMI for the full list, including what to do if you have any side effects, see Section [6. Are there any side effects?](#)

APO-IRBESARTAN HCTZ

Active ingredient(s): *irbesartan and hydrochlorothiazide*

Consumer Medicine Information (CMI)

This leaflet provides important information about taking APO-IRBESARTAN HCTZ. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using APO-IRBESARTAN HCTZ.**

Where to find information in this leaflet:

- [1. Why am I taking APO-IRBESARTAN HCTZ?](#)
- [2. What should I know before I take APO-IRBESARTAN HCTZ?](#)
- [3. What if I am taking other medicines?](#)
- [4. How do I take APO-IRBESARTAN HCTZ?](#)
- [5. What should I know while taking APO-IRBESARTAN HCTZ?](#)
- [6. Are there any side effects?](#)
- [7. Product details](#)

1. Why am I taking APO-IRBESARTAN HCTZ?

APO-IRBESARTAN HCTZ contains the active ingredients **irbesartan and hydrochlorothiazide**. Both medicines reduce blood pressure in different ways.

Irbesartan belongs to a group of medicines known as angiotensin-II receptor antagonists. Angiotensin II is a substance produced in the body that causes blood vessels to narrow. Irbesartan blocks angiotensin-II and therefore widens your blood vessels, making it easier for your heart to pump blood throughout your body. This helps to lower your blood pressure.

Hydrochlorothiazide belongs to the class of medicines known as diuretics. Diuretics cause an increase in the volume of urine. They also help with lowering blood pressure particularly when combined with other blood pressure reducing medicines.

APO-IRBESARTAN HCTZ is used to treat high blood pressure, which doctors call hypertension.

2. What should I know before I take APO-IRBESARTAN HCTZ?

Warnings

Do not take APO-IRBESARTAN HCTZ if:

- you are allergic to irbesartan, hydrochlorothiazide or to sulfonamide derived medicines (commonly known as sulfur drugs), or to any ingredients listed at the end of this leaflet. Always check the ingredients to make sure you can take this medicine.

- you are pregnant (or think you may be pregnant) or are planning to become pregnant.
- you are breastfeeding.
- you have diabetes or have kidney problems and you are being treated with medicines that lower blood pressure such as an ACE inhibitor, any aliskiren-containing medicines that belong to a group of medicines known as AIIIRAs (medicines also used to treat high blood pressure).
- you are not producing urine.

Check with your doctor if you:

- have any other medical conditions.
- have had recent excessive vomiting or diarrhoea or think you are dehydrated.
- have kidney problems, or have had a kidney transplant or dialysis.
- have heart problems.
- have liver problems, or have had liver problems in the past.
- have diabetes.
- have gout or have had gout in the past.
- have lupus erythematosus.
- have high or low levels of potassium or sodium or other electrolytes in your blood.
- have primary aldosteronism. This is a condition which causes sodium retention and an increase in blood pressure.
- are strictly restricting your salt intake.
- have had a sympathectomy. This is a surgical procedure to cut or block a nerve in your body.
- have been taking diuretics. These are medicines that reduce fluid retention in your body and are often used to treat high blood pressure.
- have had skin cancer or you develop an unexpected skin lesion during the treatment. Treatment with hydrochlorothiazide, particularly long term use with high doses, may increase the risk of some types of skin and lip cancer (non-melanoma skin cancer). Protect your skin from sun exposure and UV rays while taking APO-IRBESARTAN HCTZ.
- take any medicines for any other condition.
- are pregnant or intend to become pregnant.
- if you are breastfeeding or plan to breastfeed.
- plan to have surgery (even at the dentist) that needs a general anaesthetic.

- have a history of allergy or asthma.
- have allergies to any substances, such as foods, preservatives or dyes.
- have a decrease in your vision or pain in one or both of your eyes. If so, you should discontinue APO-IRBESARTAN HCTZ treatment and seek prompt medical attention, as these could be symptoms of fluid accumulation in the eye (choroidal effusion) or an increase of pressure in your eye (glaucoma) and can happen within hours to weeks of taking APO-IRBESARTAN HCTZ. This can lead to permanent vision loss, if not treated. If you have had a penicillin or sulfonamide (sulfur drug) allergy, you can be at higher risk of developing this.
- experienced breathing or lung problems (including inflammation or fluid in the lungs) following hydrochlorothiazide intake in the past.

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

Use in children

APO-IRBESARTAN HCTZ should not be given to children.

Pregnancy and breastfeeding

- Check with your doctor if you are pregnant or intend to become pregnant. Do not take APO-IRBESARTAN HCTZ if you are pregnant or intend to become pregnant.
- Talk to your doctor if you are breastfeeding or intend to breastfeed. Do not take APO-IRBESARTAN HCTZ if you are breastfeeding.

3. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

Tell your doctor if you are taking or plan to take any of the following:

- other medicines to treat high blood pressure.
- other diuretics.
- lithium or lithium-containing medicines (for example lithium carbonate).
- potassium tablets.
- potassium-containing salt tablets.
- non-steroidal anti-inflammatory medicine such as diclofenac, ibuprofen) and COX-2 inhibitors (for example celecoxib). These medicines may be used to relieve pain, swelling and other symptoms of inflammation including arthritis. Taking APO-IRBESARTAN HCTZ and an anti-inflammatory medicine alone or with a thiazide diuretic (fluid tablet)

may damage your kidneys. It may also reduce the effect APO-IRBESARTAN HCTZ has on lowering blood pressure.

- alcoholic drinks.
- sleeping tablets.
- strong pain medicines such as codeine or morphine.
- medicines to treat diabetes, such as repaglinide. APO-IRBESARTAN HCTZ might induce hypoglycaemia (low blood sugar).
- calcium supplements or medicines containing calcium.
- medicines containing Vitamin D.
- medicines for gout.
- powder or granules used to help reduce cholesterol (cholestyramine or colestipol hydrochloride).
- heart medicines such as digoxin or medicines to treat abnormal heart rhythm such as sotalol hydrochloride.
- corticosteroid medicines such as prednisone, cortisone or adrenocorticotrophic hormone (ACTH).
- medicines used to treat cancer.
- amantadine, a medicine used to treat Parkinson's disease or to prevent influenza.
- anticholinergic medicines, these can be used to treat Parkinson's disease, to relieve stomach cramps or spasms or used to prevent travel sickness.
- carbamazepine, a medicine used to treat convulsion.
- medicines used during surgery.
- medicines used in an emergency such as adrenaline.

Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect APO-IRBESARTAN HCTZ.

4. How do I take APO-IRBESARTAN HCTZ?

How much to take

- APO-IRBESARTAN HCTZ will usually be prescribed by your doctor if previous treatment does not produce a sufficient drop in your blood pressure. Your doctor will tell you how to switch from your previous treatment to APO-IRBESARTAN HCTZ.
- The usual dose of APO-IRBESARTAN HCTZ is one tablet once a day. The full blood pressure lowering effect should be reached 6 to 8 weeks after beginning treatment.
- If your blood pressure is not satisfactorily reduced with APO-IRBESARTAN HCTZ, your doctor may prescribe another medicine to be taken with APO-IRBESARTAN HCTZ.
- Ask your doctor or pharmacist if you are unsure of the correct dose for you. They will tell you exactly how much to take.

- Follow the instructions provided and take APO-IRBESARTAN HCTZ until your doctor tells you to stop.

When to take APO-IRBESARTAN HCTZ

- Take APO-IRBESARTAN HCTZ about the same time each day.

How to take APO-IRBESARTAN HCTZ

- Swallow the tablet whole with a glass of water. It does not matter whether you take APO-IRBESARTAN HCTZ tablets before or after food.
- Continue taking APO-IRBESARTAN HCTZ until your doctor tells you to stop.
- To help you remember to take your tablets each day, APO-IRBESARTAN HCTZ tablets are supplied in a calendar pack with the foil backing marked with the days of the week. This is just a way to help you to remember to take your tablets. All of the tablets in the pack are the same.
- When you start a new strip of tablets, take the tablet marked "START". On the next day, take the tablet marked with the relevant day of the week. Continue taking your tablets each day until all of the tablets are taken. Commence the next strip at "START" and continue as before.

If you forget to take APO-IRBESARTAN HCTZ

APO-IRBESARTAN HCTZ should be used regularly at the same time each day.

If you miss your dose at the usual time and if it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Otherwise, take it as soon as you remember, and go back to taking your tablets as you would normally.

Do not take a double dose to make up for the dose you missed.

If you take too much APO-IRBESARTAN HCTZ

If you think that you have taken too much APO-IRBESARTAN HCTZ, you may need urgent medical attention.

You should immediately:

- phone the Poisons Information Centre (**by calling 13 11 26**), or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

You should do this even if there are no signs of discomfort or poisoning.

If you take too many APO-IRBESARTAN HCTZ tablets you will probably feel light-headed or dizzy.

5. What should I know while taking APO-IRBESARTAN HCTZ?

Things you should do

- Tell any other doctors, dentists, and pharmacists who are treating you that you are taking APO-IRBESARTAN HCTZ.
- Tell your doctor immediately if you become pregnant while taking APO-IRBESARTAN HCTZ.
- Have your blood pressure checked when your doctor tells you to, to make sure APO-IRBESARTAN HCTZ is working.
- If you are about to be started on any new medicine, tell your doctor and pharmacist that you are taking APO-IRBESARTAN HCTZ.
- If you plan to have surgery that needs a general anaesthetic, tell your doctor or dentist that you are taking APO-IRBESARTAN HCTZ. Your blood pressure may drop suddenly.
- Make sure you drink enough water during exercise and hot weather when you are taking APO-IRBESARTAN HCTZ, especially if you sweat a lot. If you do not drink enough water while taking APO-IRBESARTAN HCTZ, you may faint or feel light-headed or sick. This is because your body does not have enough fluid and your blood pressure is low. If you continue to feel unwell, tell your doctor.
- Tell your doctor if you have excessive vomiting and/or diarrhoea while taking APO-IRBESARTAN HCTZ. You may lose too much water and salt and your blood pressure may drop too much.
- Tell your doctor immediately if you feel light-headed or dizzy after taking your first dose of APO-IRBESARTAN HCTZ, or when your dose is increased.
- Tell your doctor if you experience an increased sensitivity of the skin to the sun with symptoms of sunburn (such as redness, itching, swelling, blistering) occurring more quickly than normal.

- If you develop any shortness of breath or difficulty breathing after taking APO-IRBESARTAN HCTZ, seek medical attention immediately.
- Tell your doctor if you experience abdominal pain, nausea, vomiting or diarrhoea after taking APO-IRBESARTAN HCTZ. Your doctor will decide on further treatment. Do not stop taking APO-IRBESARTAN HCTZ on your own.

Things you should not do

- Do not give APO-IRBESARTAN HCTZ to anyone else, even if they have the same condition as you.
- Do not take APO-IRBESARTAN HCTZ to treat any other complaints unless your doctor or pharmacist tells you to.
- Do not stop taking APO-IRBESARTAN HCTZ or lower the dosage without checking with your doctor.

Things to be careful of

- If you feel light-headed, dizzy or faint when getting out of bed or standing up, get up slowly.
- Standing up slowly, especially when you get up from bed or chairs will help your body get used to the change in position and blood pressure. If this problem continues or gets worse, talk to your doctor.
- The hydrochlorothiazide contained in this medicine could produce a positive analytical result in an antidoping test.

Driving or using machines

Be careful before you drive or use any machines or tools until you know how APO-IRBESARTAN HCTZ affects you.

As with many other medicines used to treat high blood pressure, APO-IRBESARTAN HCTZ may cause dizziness or light-headedness in some people. If this occurs do not drive.

Make sure you know how you react to APO-IRBESARTAN HCTZ before you drive a car, operate machinery or do anything else that could be dangerous if you are dizzy or light-headed.

Drinking alcohol

Tell your doctor if you drink alcohol.

If you drink alcohol, dizziness or light-headedness may be worse.

Looking after your medicine

- Keep APO-IRBESARTAN HCTZ tablets in a cool dry place where the temperature stays below 25°C.
- Keep your tablets in the blister pack until it is time to take them.

Follow the instructions in the carton on how to take care of your medicine properly.

Store it in a cool dry place away from moisture, heat or sunlight; for example, do not store it:

- in the bathroom or near a sink, or
- in the car or on window sills.

Keep it where young children cannot reach it.

Getting rid of any unwanted medicine

If you no longer need to use this medicine or it is out of date, take it to any pharmacy for safe disposal.

Do not use this medicine after the expiry date.

6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

Less serious side effects

Less serious side effects	What to do
<ul style="list-style-type: none"> • Headache. • Dizziness or light-headedness (vertigo). • Unusual tiredness or weakness, fatigue. • Shortness of breath when exercising or looking pale. • Pain the stomach or gut. • Nausea and/or vomiting. • Sexual problems. • Low blood glucose levels. 	<p>Speak to your doctor if you have any of these less serious side effects and they worry you.</p> <p>They are generally mild and do not normally require treatment to be interrupted.</p>

Serious side effects

Serious side effects	What to do
<ul style="list-style-type: none"> • Skin rash or itchiness (this may be a symptom of an allergic reaction). • Aching muscles, not caused by exercise. • Muscle pain or weakness. • Blood tests may show raised levels of an enzyme that measures breakdown of muscle (creatinine phosphokinase). • Buzzing, ringing or other persistent noise in the ear. • Changes in heart rhythm. • You are not urinating (passing water) as much as normal. • Numbness or tingling in fingers or 	<p>Stop taking APO-irbesartan HCTZ and call your doctor straight away or go to the Emergency Department at your nearest hospital if you notice any of these serious side effects.</p> <p>These are serious side effects. You may need urgent medical attention or</p>

Serious side effects	What to do
<p>toes.</p> <ul style="list-style-type: none"> • Painful, swollen joints that may be symptoms of gout. • Fainting. • Symptoms that may indicate low platelet count such as easy excessive bruising, bleeding from gums or nose, prolonged bleeding from cuts and blood in urine or stools. • Disturbances in your vision (such as blurred vision and seeing haloes around lights) and eye pain. • Symptoms that may occur if you have developed diabetes, such as excessive thirst, passing greatly increased amounts of urine, increased appetite with weight loss, feeling tired, drowsy, weak, depressed, irritable and generally unwell. • Symptoms that may indicate kidney disease such as passing little or no urine, drowsiness, nausea, vomiting, breathlessness, loss of appetite and weakness. • Symptoms that may indicate high potassium levels in the blood such as nausea, diarrhoea, muscle weakness and changes in heart rhythm. • Yellowing of the skin and/or eyes, also known as jaundice. • Symptoms that may indicate liver disease such as nausea, vomiting, loss of appetite, feeling generally unwell, fever, itching, yellowing of the skin and eyes and dark coloured urine. • Swelling of the face, lips, tongue or throat which may cause difficulty in swallowing or breathing. • Severe and sudden onset of pinkish, itchy swellings on the skin, also called hives or nettle rash. • Acute respiratory distress (signs include shortness of breath, fever, weakness and confusion). 	<p>hospitalisation. These side effects are very rare.</p>

Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people.

Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at www.tga.gov.au/reporting-problems. By reporting side effects, you can help provide more information on the safety of this medicine.

Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.

7. Product details

This medicine is only available with a doctor's prescription.

What APO-IRBESARTAN HCTZ contains

Active ingredient (main ingredient)	<ul style="list-style-type: none"> • Each tablet contains either 150 mg or 300 mg of irbesartan and 12.5 mg or 25 mg of hydrochlorothiazide as the active ingredients.
Other ingredients (inactive ingredients)	<ul style="list-style-type: none"> • Croscarmellose sodium • Lactose monohydrate • Magnesium stearate • Pigment Blend PB-24899 PINK • Povidone • 150/12.5 and 300/12.5 tablets also contain OPADRY II complete film coating system 30F84418 PINK. • 300/25 tablets also contain OPADRY II Complete film coating system 30F86974 BROWN.
Potential allergens	This medicine contains sugars as lactose.

This medicine does not contain gluten, tartrazine or any other azo dyes.

Do not take this medicine if you are allergic to any of these ingredients.

What APO-IRBESARTAN HCTZ looks like

150/12.5 mg tablets*

Peach coloured, oval shaped, biconvex, film coated tablet, debossed with "L183" on one side and plain on the other side.

AUST R 213302.

300/12.5 mg tablets*

Peach coloured, oval shaped, biconvex, film coated tablet, debossed with "L184" on one side and plain on the other side.

AUST R 213303.

300/25 mg tablets*

Pinkish brown, oval shaped, biconvex, film coated tablet, debossed with "L185" on one side and plain on the other side.

AUST R 213305.

Available in blister packs of 30 tablets.

* Not all strengths may be available.

Who distributes APO-IRBESARTAN HCTZ

Arrotex Pharmaceuticals Pty Ltd

15-17 Chapel St

Cremorne VIC 311

This leaflet was prepared in August 2025.