

ORATANE®

Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

1. Why am I using ORATANE®?

ORATANE® contains the active ingredient isotretinoin. ORATANE® is used to treat acne. For more information, see Section [1. Why am I using ORATANE®?](#) in the full CMI.

2. What should I know before I use ORATANE®?

Do not use if you have ever had an allergic reaction to ORATANE® or any of the ingredients listed at the end of the CMI. **Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.** For more information, see Section [2. What should I know before I use ORATANE®?](#) in the full CMI.

3. What if I am taking other medicines?

Some medicines may interfere with ORATANE® and affect how it works. A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

4. How do I use ORATANE®?

- Your doctor will tell you how many ORATANE® capsules to take each day, take them exactly as your doctor has prescribed.
- ORATANE® capsules should be swallowed whole with a glass of water or milk and must always be taken with food.

More instructions can be found in Section [4. How do I use ORATANE®?](#) in the full CMI.

5. What should I know while using ORATANE®?

Things you should do	<ul style="list-style-type: none">• Remind any doctor, dentist or pharmacist you visit that you are using ORATANE®.• Use effective contraception for one month before, during and one month after treatment.• If you become pregnant while taking ORATANE®, stop taking it and tell your doctor immediately.• Tell your doctor if you are intending to do a lot of heavy lifting or exercise.• Your skin may be more sensitive while on ORATANE®. Avoid excessive sun exposure and certain skin and hair treatments.
Things you should not do	<ul style="list-style-type: none">• Do not stop taking ORATANE® suddenly, change the dose or let yourself run out of medicine over the weekend or on holidays.• Do not give ORATANE® to anyone else even if their symptoms seem similar to yours.• Do not use ORATANE® to treat other complaints unless your doctor says to.• Do not donate blood during treatment with ORATANE® or for at least 1 month after stopping treatment.
Driving or using machines	<ul style="list-style-type: none">• Be careful driving or operating machinery until you know how ORATANE® affects you.
Drinking alcohol	<ul style="list-style-type: none">• You must tell your doctor if you drink large amounts of alcohol.
Looking after your medicine	<ul style="list-style-type: none">• Keep the blister pack in a cool dry place where the temperature stays below 25°C.• Protect ORATANE® from light.

For more information, see Section [5. What should I know while using ORATANE®?](#) in the full CMI.

6. Are there any side effects?

ORATANE® helps most people with acne but it may have unwanted side effects in a few people. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects. When you are using ORATANE®, you can have some serious side effects. Serious side effects may include nausea, vomiting, persistent headache, blurred vision or visual disturbances, changes in your hearing or ringing in your ears, severe upper stomach pain, unexpected muscle pain, tenderness or weakness, blood in stools or severe diarrhoea, severe bruising, sudden red or itchy spots, painful red areas, fever and chills, aching muscles, feeling depressed with or without suicidal thoughts and thinking, seeing or hearing things that are not real.

For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

ORATANE[®] (orr·a·tayn)

Active ingredient(s): *isotretinoin*

Consumer Medicine Information (CMI)

This leaflet provides important information about using ORATANE[®]. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using ORATANE[®].**

Where to find information in this leaflet:

1. [Why am I using ORATANE[®]?](#)
2. [What should I know before I use ORATANE[®]?](#)
3. [What if I am taking other medicines?](#)
4. [How do I use ORATANE[®]?](#)
5. [What should I know while using ORATANE[®]?](#)
6. [Are there any side effects?](#)
7. [Product details](#)

1. Why am I using ORATANE[®]?

ORATANE[®] contains the active ingredient isotretinoin.

ORATANE[®] belongs to a group of medicines called retinoids, which are similar to vitamin A.

Retinoids work by reducing the amount of the oily substance (i.e. sebum) made by glands in your skin, reducing bacteria and inflammation and opening clogged pores.

ORATANE[®] is used to treat acne. There are many different types of medicines used to treat acne.

ORATANE[®] is used for more severe cases.

Your doctor, however, may have prescribed ORATANE[®] for another purpose.

Ask your doctor if you have any questions about why ORATANE[®] has been prescribed for you.

This medicine is available only with a doctor's prescription. ORATANE[®] is not addictive.

2. What should I know before I use ORATANE[®]?

Warnings

Do not use ORATANE[®] if:

- you are allergic to isotretinoin, have had an allergic reaction to ORATANE[®], vitamin A, other retinoids or any of the ingredients listed at the end of this leaflet. Always check the ingredients to make sure you can use this medicine.
- you are pregnant, or for at least one month before you plan to fall pregnant. If you fall pregnant while taking ORATANE[®] there is an extremely high risk of having a baby that is severely deformed. You must use effective contraception for one month before, during and one month after treatment.
- you are breastfeeding. You must stop breastfeeding before treatment begins. Do not breastfeed while taking ORATANE[®].
- you are taking tetracycline antibiotics such

asdoxycycline

- you have severe liver disease
- you have very high fat levels (cholesterol, triglycerides) in your blood
- you have hypervitaminosis A. This is a condition caused by an excessive amount of vitamin A in the diet.
- depression (patients should have a mental health assessment before starting treatment)
- difficulty getting or maintaining an erection (in males)
- reduced sex drive or ability to feel pleasure
- vaginal dryness (in females)
- breast swelling in males
- the packaging is torn or shows signs of tampering
- the expiry date (EXP) printed on the pack has passed. If you take this medicine after the expiry date has passed, it may not work as well.

If you are not sure if you should start taking ORATANE[®], contact your doctor.

Do not give ORATANE[®] to children.

ORATANE[®] may be associated with slowing of growth when used in children before puberty. The use of ORATANE[®] in children less than 12 years of age is not recommended.

Check with your doctor if you:

- you drink large amounts of alcohol
- you have any allergies to any other medicines, foods, preservatives or dyes. ORATANE[®] capsules contain soya oil, which may contain traces of arachidic acid (a component of peanut oil).
- you have or have had any other health problems or issues including:
 - diabetes, or a history of diabetes in your family
 - depression
 - liver disease
 - kidney disease
 - lipid (cholesterol or triglyceride) disorder
 - hormone disorder
 - eye disorders
 - stomach or bowel disease

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

Pregnancy and breastfeeding

If you become pregnant while taking ORATANE[®], stop taking it and tell your doctor immediately.

ORATANE[®] can cause birth defects (damage to unborn babies). You must use strict birth control, starting at least 1 month before you begin taking ORATANE[®], for the whole time you are taking ORATANE[®] and for 1 month after you finish taking ORATANE[®].

There is no known risk to males who wish to father children.

Talk to your doctor if you are breastfeeding or intend to breastfeed.

3. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines may interfere with ORATANE® and affect how it works. These include:

- tetracycline antibiotics such as Doxycycline
- vitamin A, or preparations containing vitamin A (including vitamin supplements)
- other medicines you are using to treat your acne
- the "mini-pill", a progestogen-only oral contraceptive pill

These medicines may be affected by ORATANE® or may affect how well it works. You may need to take different amounts of your medicine, or you may need to take different medicines. Your doctor will advise you.

Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect ORATANE®.

4. How do I use ORATANE®?

How much to take

- Follow all directions given to you by your doctor or pharmacist carefully.
- They may differ from the information contained in this leaflet.
- Take ORATANE® exactly as your doctor has prescribed.
- Your doctor will tell you how many ORATANE® capsules to take each day.
- The dose will be calculated to suit your individual needs and your body weight. This dose may be adjusted during treatment when your doctor knows how you respond to ORATANE®.

When to take ORATANE®

- ORATANE® may be taken once or twice a day and must always be taken with food.
- Female patients should wait until the 2nd or 3rd day of the next normal menstrual period before starting ORATANE® treatment. This helps ensure that you aren't pregnant before you start taking ORATANE®.

How to take ORATANE®

- Capsules should be swallowed whole with a glass of water or milk.
- Do not open the capsules and do not take any capsules that are damaged.

How long to take ORATANE®

- Continue taking ORATANE® for as long as your doctor prescribes.
- Acne treatment with ORATANE® will usually last 4 to 8 months. In the first few weeks of treatment your acne may get a little worse before it gets better. This is a sign that ORATANE® is working.
- At the end of this time your acne should have cleared up significantly. Most patients notice their skin condition continues to improve even after ORATANE® treatment is finished.
- Please note that ORATANE® cannot improve scars or pitting that were present before treatment started, but it will help prevent such skin damage in the future.

If you forget to use ORATANE®

ORATANE® should be used regularly at the same time each day.

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Do not take a double dose to make up for the dose you missed.

- This may increase the chance of getting an unwanted side effect.
- If you have missed several doses, please inform your doctor and follow the advice given to you.

If you take too much ORATANE® (overdose)

If you think that you have used too much ORATANE®, you may need urgent medical attention.

You should immediately:

- phone the Poisons Information Centre (by calling 13 11 26), or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

You should do this even if there are no signs of discomfort or poisoning.

You may need urgent medical attention.

Signs of overdose include transient headache; vomiting; facial flushing; reddened, cracked lips; stomach pain; headache; dizziness and unsteady walking.

5. What should I know while using ORATANE®?

Call your doctor straight away if you become pregnant while taking ORATANE®, stop taking it and tell your doctor immediately.

Things you should do

- Remind any doctor, dentist or pharmacist you visit that you are using ORATANE®.
- Tell your doctor if, for any reason, you have not taken your medicine exactly as prescribed. Otherwise, your doctor may think that it was not effective and change your treatment unnecessarily.

- Tell your doctor if you feel ORATANE® capsules are not helping your condition.
- Be sure to keep all of your appointments with your doctor so that your progress can be checked. Your doctor may ask you to have regular blood tests to monitor your liver function, blood sugar levels and blood cholesterol levels.
- If you are intending to do a lot of heavy lifting or exercise, tell your doctor. Your muscles and joints may be more prone to tenderness or stiffness if you do a lot of heavy exercise while taking ORATANE®.
- If you experience dryness of the lips, mouth, nose or skin, you can use a moisturiser or petroleum jelly to soften the areas not affected by the acne.
- Talk to your doctor if you experience persistent pain in your lower back or buttocks during treatment with ORATANE®. These symptoms may be signs of sacroiliitis, a type of inflammatory back pain. Your doctor may discontinue treatment with ORATANE® and refer you to a specialist for treatment of inflammatory back pain. Further evaluation may be needed including imaging modalities such as MRI.

Things you should not do

- Do not stop taking ORATANE® suddenly or change the dose without first checking with your doctor.
- Do not let yourself run out of medicine over the weekend or on holidays.
- Do not give ORATANE® to anyone else even if their symptoms seem similar to yours.
- Do not use ORATANE® to treat other complaints unless your doctor says to.
- Do not donate blood during treatment with ORATANE® or for at least 1 month after stopping treatment.

Things to be careful of

- Wearing contact lenses during treatment with ORATANE® may cause discomfort. ORATANE® may cause dry eyes. An eye lubricant or artificial tears, available from your pharmacist, should relieve this problem. Otherwise, you may temporarily need to wear your lenses for shorter periods or wear glasses instead.
- Avoid excessive sun exposure and solariums and apply sunscreen whilst taking ORATANE®. Your skin may be more prone to sunburn while on ORATANE®.
- Avoid waxing and dermabrasion whilst taking ORATANE® and for 5 to 6 months after stopping ORATANE® treatment. Your skin may be more sensitive while on ORATANE®. Waxing may cause dermatitis and dermabrasion may cause scarring during and for several months after ORATANE® treatment.
- Avoid using facial peels, electrolysis and some hair treatments. Your skin and hair may be more delicate during treatment and for a while after ORATANE® treatment.

Driving or using machines

Be careful before you drive or use any machines or tools until you know how ORATANE® affects you.

Normally ORATANE® would not affect your ability to drive a car or operate machinery. However, altered night vision and other visual disturbances may occur when taking ORATANE®. Make sure you know how you react to ORATANE® before you drive a car, operate machinery or do anything else that may be dangerous if your vision is affected.

Drinking alcohol

Tell your doctor if you drink alcohol.

Looking after your medicine

- Keep your capsules in the blister pack until it is time to take them. If you take the capsules out of the packaging, they will not keep as well.
- Store it in a cool dry place where the temperature stays below 25°C for ORATANE® 5mg, 10mg, 30mg & 40mg and , below 30°C for ORATANE® 20mg, away from moisture, heat or sunlight; for example, do not store it:
 - in the bathroom or near a sink, or
 - in the car or on windowsills.

Keep ORATANE® where young children cannot reach it.

When to discard your medicine

If your doctor tells you to stop taking ORATANE®, or the capsules have passed their expiry date, ask your pharmacist what to do with any capsules that are left over.

Getting rid of any unwanted medicine

If you no longer need to use this medicine or it is out of date, take it to any pharmacy for safe disposal.

Do not use this medicine after the expiry date.

6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

Less serious side effects

Less serious side effects	What to do
<p>Symptoms associated with an excess of Vitamin A in the body (hypervitaminosis A):</p> <ul style="list-style-type: none"> • dryness of the lips, mouth, nose and skin • peeling palms of the hands and soles of the feet • eye problems such as dry, sore, swollen or itchy eyes, discharge or trouble seeing at night 	<p>Speak to your doctor if you have any of these less serious side effects and they worry you.</p>

Less serious side effects	What to do
<ul style="list-style-type: none"> hoarseness nosebleeds <p>Symptoms associated with skin and appendage disorders:</p> <ul style="list-style-type: none"> fragile skin change in colour of the skin itchy skin rash an increased susceptibility to sunburn flaring of acne, usually at the start of treatment sweating changes to the nails hair loss (sometimes occurs and is usually temporary but in rare cases has persisted) excessive hairiness <p>Symptoms associated with musculoskeletal system disorders:</p> <ul style="list-style-type: none"> tenderness or stiffness in your bones, joints or muscles muscle or joint pain <p>Symptoms associated with central nervous system:</p> <ul style="list-style-type: none"> headache tiredness <p>Symptoms associated with renal and urinary disorders:</p> <ul style="list-style-type: none"> inflammation of the urethra 	

Serious side effects	What to do
<p>Symptoms associated with skin and appendage disorders:</p> <ul style="list-style-type: none"> sudden red, often itchy spots, similar to the rash of measles starting on the face, hands or feet. The spots may blister or change to flat round raised, red, pale-centered marks. You may also have a fever, sore throat, headache and/ or diarrhoea. painful red areas, that change to large blisters and end with peeling of layers of skin, that may occur on lips, mouth, eyes, nose and genitals. Those affected may have fever and chills, aching muscles and generally feel unwell. <p>Symptoms associated with the gastrointestinal system:</p> <ul style="list-style-type: none"> nausea vomiting severe upper stomach pain 	<p>Call your doctor straight away or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.</p>

Serious side effects	What to do
<ul style="list-style-type: none"> blood in stools or severe diarrhoea <p>Symptoms associated with the central nervous system:</p> <ul style="list-style-type: none"> persistent headache <p>Symptoms associated with sensory disorders:</p> <ul style="list-style-type: none"> blurred vision or visual disturbances changes in your hearing or ringing in your ears thinking, seeing or hearing things that are not real <p>Symptoms associated with musculoskeletal system disorders:</p> <ul style="list-style-type: none"> unexpected muscle pain, tenderness or weakness severe bruising persistent pain in the lower back or buttocks <p>Symptoms associated with depression:</p> <ul style="list-style-type: none"> feeling depressed, with or without suicidal thoughts feeling sad or having crying spells losing interest in activities you once enjoyed sleeping too much or having trouble sleeping changes in your appetite or body weight having trouble concentrating withdrawing from your friends or family feeling like you have no energy feelings of worthlessness or inappropriate guilt 	

Serious side effects

Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people.

Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at www.tga.gov.au/reporting-problems. By reporting side effects, you can help provide more information on the safety of this medicine.

Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.

7. Product details

This medicine is only available with an authorised doctor's prescription.

What ORATANE® contains

Active ingredient (main ingredient)	isotretinoin
Other ingredients (inactive ingredients)	<p>Capsule Filling</p> <ul style="list-style-type: none"> • Beeswax – yellow • Disodium edetate • Butylated Hydroxyanisole • dl-alpha-tocopherol • Soya oil – partially hydrogenated • Soya oil • Vegetable oil – hydrogenated (5mg, 10mg and 20mg only) • Soya oil – hydrogenated (30mg and 40mg only). <p>Capsule Shell</p> <ul style="list-style-type: none"> • Sorbitol solution (70% non-crystallising) • Glycerol • Gelatin† • Titanium dioxide • Water – purified • Brilliant scarlet 4R CI16255 (10mg and 20mg only) • Sunset yellow FCF CI15985 (40mg only) • Indigo carmine CI73015 (20mg only) • Iron oxide black CI177499 (10mg only) • Iron oxide red (30mg only)
Potential allergens	<p>Soya oil may contain traces of arachidic acid (a component of peanut oil)</p> <p>†Contains sulfites. May contain trace amounts of phenylalanine.</p>

Do not take this medicine if you are allergic to any of these ingredients.

ORATANE® is available as a 20 mg capsule in a pack of 60 capsules.

What ORATANE® looks like:

ORATANE® 5 mg capsules are soft, oval, faint pinkish/cream to cream in colour containing a yellow/orange liquid. Available in blister packs of 60 capsules. ARTG 127499

ORATANE® 10 mg capsules are soft, oval, opaque light violet in colour containing a yellow/orange liquid. Available in blister packs of 60 capsules. ARTG 75034

ORATANE® 20 mg capsules are soft, oval, opaque maroon, red in colour containing a yellow/orange liquid. Available in blister packs of 60 capsules. ARTG 69868

ORATANE® 30 mg capsules are pink coloured soft gelatin capsules containing a yellow/ orange liquid. Available in blister pack of 60 capsules. ARTG 303071

ORATANE® 40 mg capsules are soft, oval, opaque orange in colour containing a yellow/orange liquid. Available in blister packs of 30 capsules. ARTG 117206

Sponsor:

Arrotex Pharmaceuticals Pty Ltd
15 – 17 Chapel Street
Cremorne Victoria 3121
www.arrotex.com.au

This leaflet was prepared in March 2025