

# ROCURONIUM BAXTER

## Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

### 1. Why am I using ROCURONIUM BAXTER?

ROCURONIUM BAXTER contains the active ingredient rocuronium bromide. ROCURONIUM BAXTER is one of a group of medicines called muscle relaxants. ROCURONIUM BAXTER is used during an operation as part of the general anaesthetic. When you have an operation, your muscles must be completely relaxed. This makes it easier for the surgeon to perform the operation. For more information, see Section [1. Why am I using ROCURONIUM BAXTER?](#) in the full CMI.

### 2. What should I know before I use ROCURONIUM BAXTER

Do not use if you have ever had an allergic reaction to ROCURONIUM BAXTER or any of the ingredients listed at the end of the CMI.

**Talk to your doctor if you have any other medical conditions, take any other medicines, or are pregnant or plan to become pregnant or are breastfeeding.**

For more information, see Section [2. What should I know before I use ROCURONIUM BAXTER?](#) in the full CMI.

### 3. What if I am taking other medicines?

Some medicines may interfere with ROCURONIUM BAXTER and affect how it works.

A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

### 4. How ROCURONIUM BAXTER is given

- ROCURONIUM BAXTER will be given by a doctor. It will not be given to you until you are asleep from the anaesthetic.
  - It will be injected into a vein before and/or during an operation. It will be given as a single injection or continuous infusion.
- More instructions can be found in Section [4. How do I use ROCURONIUM BAXTER?](#) in the full CMI.

### 5. What should I know after having ROCURONIUM BAXTER?

<b>Driving or using machines</b>	<ul style="list-style-type: none"><li>• Your doctor will tell you when it is safe to drive and operate potentially dangerous machinery after you have been given ROCURONIUM BAXTER.</li></ul>
----------------------------------	---

For more information, see Section [5. What should I know after having ROCURONIUM BAXTER?](#) in the full CMI.

### 6. Are there any side effects?

Like all medicines, ROCURONIUM BAXTER can cause side effects, although not everybody gets them. Your doctor will discuss these with you and will explain the risks and benefits of using ROCURONIUM BAXTER. When you get ROCURONIUM BAXTER, you can have some serious side effects.

For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

# ROCURONIUM BAXTER

Active ingredient(s): rocuronium bromide

## Consumer Medicine Information (CMI)

This leaflet provides important information about using ROCURONIUM BAXTER. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using ROCURONIUM BAXTER.**

Where to find information in this leaflet:

- [1. Why am I using ROCURONIUM BAXTER?](#)
- [2. What should I know before I use ROCURONIUM BAXTER?](#)
- [3. What if I am taking other medicines?](#)
- [4. How ROCURONIUM BAXTER is given](#)
- [5. What should I know after having ROCURONIUM BAXTER?](#)
- [6. Are there any side effects?](#)
- [7. Product details](#)

## 1. Why am I using ROCURONIUM BAXTER?

**ROCURONIUM BAXTER contains the active ingredient rocuronium bromide.** ROCURONIUM BAXTER is one of a group of medicines called muscle relaxants.

Muscle relaxants are used during an operation as part of the general anaesthetic. When you have an operation, your muscles must be completely relaxed. This makes it easier for the surgeon to perform the operation.

Normally the nerves send messages to the muscles by impulses. ROCURONIUM BAXTER acts by blocking these impulses so the muscles are relaxed. Because the muscles needed for breathing also become relaxed you will need help with your breathing (artificial respiration) during and after your operation until you can breathe on your own.

During the operation the effect of the muscle relaxant is constantly checked and if necessary some more drug is given. At the end of the operation the effects of ROCURONIUM BAXTER are allowed to wear off and you can start breathing on your own. Sometimes another drug is given to help speed this up. ROCURONIUM BAXTER can also be used in Intensive Care to keep your muscles relaxed.

## 2. What should I know before I use ROCURONIUM BAXTER?

### Warnings

**Do not use ROCURONIUM BAXTER if:**

- you are allergic to rocuronium bromide, or any of the ingredients listed at the end of this leaflet.

- Always check the ingredients to make sure you can use this medicine.

Some of the symptoms of an allergic reaction may include shortness of breath, wheezing or difficulty breathing; swelling of the face, lips, tongue or other parts of the body; rash, itching or hives on the skin.

**ROCURONIUM BAXTER should not be given to a child under the age of one month.** The safety of administration of ROCURONIUM BAXTER has not been established in children under the age of one month.

### Before you are given it:

If you are going to have an operation it is important that you discuss the following points with your doctor, since it can influence the way ROCURONIUM BAXTER is given to you.

**Tell your doctor if you have or have had any medical conditions, especially the following:**

- an allergy to muscle relaxants
- kidney disease
- liver or gallbladder disease
- a heart disease
- diseases affecting nerves or muscles
- oedema (local or generalised swelling due to fluid)

**Certain medical conditions may affect how ROCURONIUM BAXTER works:**

- low potassium levels in the blood
- high magnesium levels in the blood
- low calcium levels in the blood
- low levels of protein in the blood
- dehydration
- too much acid in the blood
- too much carbon dioxide in the blood
- general ill-health
- overweight
- burns

If you are suffering from any of these conditions your doctor will take this into account when deciding the correct dose of ROCURONIUM BAXTER for you.

**Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.**

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and

how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

### Children and Elderly

ROCURONIUM BAXTER can be used in children (30 days to adolescence) and elderly but your doctor should first assess your medical history.

Elderly (65 years and older) may be at an increased risk of residual paralysis.

### Pregnancy and breastfeeding

Check with your doctor if you are pregnant or intend to become pregnant.

Talk to your doctor if you are breastfeeding or intend to breastfeed.

**If you have not told your doctor about any of the above, tell them before you are given ROCURONIUM BAXTER.**

## 3. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

**Some medicines may be affected by ROCURONIUM BAXTER, or affect how well it works. These include:**

- anaesthetics, medicines to make you sleep during surgery
- long term concurrent use of corticosteroids (anti-inflammatory medicines) and ROCURONIUM BAXTER in the Intensive Care Unit
- antibiotics
- lithium, a medicine used to treat bipolar disorder
- medicines used to treat heart disease or high blood pressure (quinidine, calcium channel blockers, beta-blockers and diuretics (fluid tablets))
- quinine, a medicine used to treat malaria
- magnesium salts
- lignocaine and bupivacaine (local anaesthetics)
- other muscle relaxants
- carbamazepine and phenytoin, medicines used to treat epilepsy

You may need to use different amounts of your medicines or take different medicines. Your doctor will advise you.

If you are taking magnesium sulphate to treat toxemia of pregnancy (preeclampsia), tell your doctor as the dose of ROCURONIUM BAXTER may need to be reduced.

Your doctor will have a complete list of medicines that may cause problems when used with ROCURONIUM BAXTER

**Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect ROCURONIUM BAXTER.**

## 4. How ROCURONIUM BAXTER is given

ROCURONIUM BAXTER will be given by a doctor. It will not be given to you until you are asleep from the anaesthetic.

It will be injected into a vein before and/or during an operation. It will be given as a single injection or continuous infusion.

### How much will be given

The usual dose is 0.6 mg rocuronium bromide per kg body weight and the effect lasts 30-40 minutes. During the operation your doctor will check whether ROCURONIUM BAXTER is still working. You may be given additional doses if they are needed.

### If you are given too much ROCURONIUM BAXTER

As ROCURONIUM BAXTER doses are carefully worked out and are given by a doctor experienced in its use, it is extremely unlikely that you will be given too much ROCURONIUM BAXTER. However, if this does happen, your doctor will make sure that you continue breathing artificially until you can breathe on your own again. Your doctor may speed-up your recovery by giving you a drug that reverses the effects of ROCURONIUM BAXTER.

## 5. What should I know after having ROCURONIUM BAXTER?

### Driving or using machines

**Be careful before you drive or use any machines or tools until you know how ROCURONIUM BAXTER affects you.**

Your doctor will tell you when it is safe to drive and operate potentially dangerous machinery after you have been given ROCURONIUM BAXTER.

### Looking after your medicine

ROCURONIUM BAXTER is stored in the hospital.

## 6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

### Less serious side effects

Less serious side effects	What to do
<ul style="list-style-type: none"><li>• flushing</li><li>• pain at injection site</li><li>• irritation at injection site</li><li>• red skin rash or itchy rash</li></ul>	<b>Speak to your doctor if you have any of these less serious side effects and they worry you.</b>

## Serious side effects

Serious side effects	What to do
<p><b>Signs and symptoms of cardiovascular problems:</b></p> <ul style="list-style-type: none"><li>fast heart beat</li><li>dizziness, light-headedness (low blood pressure)</li></ul> <p><b>Signs and symptoms of muscle problems:</b></p> <ul style="list-style-type: none"><li>muscle weakness or paralysis</li><li>aching muscles or weakness, not caused by exercise</li></ul> <p><b>Signs and symptoms of lung problems:</b></p> <ul style="list-style-type: none"><li>wheezing, coughing</li><li>difficulty breathing</li></ul> <p><b>Signs and symptoms of allergic reactions:</b></p> <ul style="list-style-type: none"><li>rapid, shallow breathing, cold, clammy skin, a rapid, weak pulse, dizziness, weakness and fainting</li><li>swelling of the face, lips, mouth, tongue or throat which may cause difficulty in swallowing or breathing</li><li>sudden signs of allergy such as rash, itching, hives on the skin, swelling of the face, lips, tongue or other parts of the body, shortness of breath, wheezing or trouble breathing</li><li>severe allergic coronary blood vessels spasm (Kounis syndrome) resulting in chest pain (angina) or heart attack (myocardial infarction)</li></ul> <p><b>Signs and symptoms of problems in other organs:</b></p> <ul style="list-style-type: none"><li>sudden fever with rapid heartbeat, rapid breathing and stiffness, pain and/or weakness in your muscles</li><li>Dilated pupils (mydriasis) or fixed pupils</li></ul>	<p><b>Call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice any of these serious side effects.</b></p>

**Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.**

Other side effects not listed here may occur in some people.

## Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at [www.tga.gov.au/reporting-problems](http://www.tga.gov.au/reporting-problems). By reporting side effects, you can help provide more information on the safety of this medicine.

**Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.**

## 7. Product details

This medicine is only available with a doctor's prescription.

### What ROCURONIUM BAXTER contains

<b>Active ingredient (main ingredient)</b>	Rocuronium bromide
<b>Other ingredients (inactive ingredients)</b>	Sodium acetate sodium chloride glacial acetic acid sodium hydroxide water for injections

**Do not take this medicine if you are allergic to any of these ingredients.**

### What ROCURONIUM BAXTER looks like

ROCURONIUM BAXTER is a clear, colourless to yellow or orange solution in a vial.

ROCURONIUM BAXTER 50 mg/5 mL injection is available in packs of 10 vials (AUST R 169601).

ROCURONIUM BAXTER 100 mg/10 mL injection is available in packs of 10 vials (AUST R 169603).

### Who distributes ROCURONIUM BAXTER

Baxter Healthcare Pty Ltd

1 Baxter Drive

Old Toongabbie

NSW 2146

This leaflet was prepared in January 2023.